

# PREVENTIVE CARE FOR AGES BIRTH TO 23 MONTHS

*Preventive Services for Children*

## Schedule of Visits:

Within the first 4 weeks;  
then at 2, 4, 6, 9, 12, 15 & 18 mos.

Screening	Risk Assessment & Counseling	Immunizations
<ul style="list-style-type: none"> <li>◆ <b>Every Visit</b> <ul style="list-style-type: none"> <li>✓ Head Circumference, Height Weight</li> <li>✓ Physical Exam</li> <li>✓ Developmental and behavioral assessment</li> <li>✓ Eye evaluation/objective<sup>1</sup></li> <li>✓ Hearing/subjective</li> </ul> </li> <li>◆ <b>Additional</b> <ul style="list-style-type: none"> <li>✓ Hemoglobin/Hematocrit <i>Once at 9-12 months</i></li> <li>✓ Blood lead levels <i>Starting at 9 mos.</i></li> <li>✓ PPD <i>At 12 mos. (if at risk)<sup>2</sup></i></li> <li>✓ <i>Hepatitis C (if at risk)<sup>3</sup></i></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>◆ <b>Nutrition</b> <ul style="list-style-type: none"> <li>✓ Breast Feeding</li> <li>✓ Iron-rich diet</li> <li>✓ Bottle weaning</li> </ul> </li> <li>◆ <b>Injury Prevention<sup>4</sup></b> <ul style="list-style-type: none"> <li>✓ Car seats</li> <li>✓ Poison awareness</li> <li>✓ Falls</li> <li>✓ Fire Safety</li> <li>✓ Firearms</li> </ul> </li> <li>◆ <b>Dental Health</b> <ul style="list-style-type: none"> <li>✓ Bottle use</li> </ul> </li> <li>◆ <b>Exposure to Tobacco</b></li> <li>◆ <b>Family Adjustment</b></li> <li>◆ <b>Crib Safety</b> <ul style="list-style-type: none"> <li>✓ Sleep position</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>◆ <b>All Children</b> <ul style="list-style-type: none"> <li>✓ Hepatitis B <i>At birth, 1-4 mos. and 6-18 mos.<sup>5</sup></i></li> <li>✓ DTaP <i>At 2, 4, 6 and 15-18 mos.<sup>6</sup></i></li> <li>✓ Poliovirus <i>At 2, 4, and 6-18 mos.<sup>7</sup></i></li> <li>✓ H. Influenza B <i>At 2, 4, 6 and 12-15 mos.<sup>8</sup></i></li> <li>✓ MMR <i>At 12-15 mos.<sup>9</sup></i></li> <li>✓ Varicella vaccine <i>At 12-18 mos.<sup>10</sup></i></li> <li>✓ Pneumococcal Conjugate (PCV7)<sup>11</sup></li> <li>✓ Influenza vaccine <i>Annually<sup>12</sup></i></li> <li>✓ Rotovirus Vaccine- <i>Give 3 doses- 2, 4 and 6 months of age</i></li> <li>✓ Hepatitis A- <i>2 doses- 6 months apart to children 12-23 months<sup>21</sup></i></li> <li>✓ MMRV- <i>Give 1 dose at 12-15 mos. when used in place of separate vaccines.</i></li> </ul> </li> <li>◆ <b>High Risk Children</b> <ul style="list-style-type: none"> <li>✓ <i>Palivizumab or RSV-IGIV<sup>13</sup></i></li> </ul> </li> </ul> <p><i>See attached immunization schedule</i></p>