

Exondys 51TM (eteplirsen)

(Intravenous)

Effective Date: 12/01/2019

Review Date: 11/27/2019, 1/29/2020, 04/29/2021, 2/24/2022, 8/25/2022, 2/1/2023, 9/14/2023,

12/07/2023, 01/04/2024, 08/28/2024, 05/07/2025

Scope: Medicaid, Commercial, Medicare

I. Length of Authorization

Coverage will be for 6 months and may be renewed.

II. Dosing Limits

A. Max Units (per dose and over time) [HCPCS Unit]:

• 1400 billable units every 28 days

III. Summary of Evidence

Exondys 51 is an antisense oligonucleotide indicated for the treatment of Duchenne muscular dystrophy (DMD) in patients who have a confirmed mutation of the DMD gene that is amenable to exon 51 skipping. Clinical trials have demonstrated an increase in dystrophin in skeletal muscle in patients who have a confirmed mutation of the DMD gene that is amenable to exon 51 skipping. Adverse events commonly reporting in clinical trials include balance disorder and vomiting.

IV. Initial Approval Criteria 1-10

Coverage is provided in the following conditions:

Universal Criteria

Medicare members who have previously received this medication within the past 365 days are not subject to Step Therapy Requirements.

- Patient is not on concomitant therapy with other DMD-directed antisense oligonucleotides (e.g., Vyondys 53 (golodirsen), Amondys 45 (casimersen), Viltepso (viltolarsen), etc.); AND
- Patient has never received and will not receive therapy with Elevidys (delandistrogene moxeparvovec-rokl) within 6 months of this request; AND

Duchenne Muscular Dystrophy (DMD) † Φ

- Patient must have a confirmed mutation of the DMD gene that is amenable to exon 51 skipping;
 AND
- Patient has been on a stable dose of corticosteroids, unless contraindicated or intolerant, for at least 6 months; **AND**
- Patient retains meaningful voluntary motor function (e.g., patient is able to speak, manipulate objects using upper extremities, ambulate, etc.); **AND**
- Patient is receiving physical and/or occupational therapy; **AND**
- Baseline documentation of one or more of the following:
 - o Dystrophin level
 - O Timed function tests (e.g., 6-minute walk test (6MWT) time to stand [TTSTAND], time to run/walk 10 meters [TTRW], time to climb 4 stairs [TTCLIMB], etc.)
 - o Upper limb function (ULM) test
 - o North Star Ambulatory Assessment (NSAA) score
 - o Forced Vital Capacity (FVC) percent predicted

† FDA-labeled indication(s); ‡ Compendia recommended indication(s); ♠ Orphan Drug

V. Renewal Criteria 1

- Patient continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section IV; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: hypersensitivity reactions (e.g., bronchospasm, chest pain, cough, tachycardia, urticaria, etc.), etc.; **AND**
- Patient has responded to therapy compared to pretreatment baseline in one or more of the following (not all-inclusive):
 - Increase in dystrophin level
 - o Improvement in quality of life
 - O Stability, improvement, or slowed rate of decline in one of the following:

- Timed function tests (e.g., 6-minute walk test [6MWT], time to stand [TTSTAND], time to run/walk 10 meters [TTRW], time to climb 4 stairs [TTCLIMB])
- Upper limb function (ULM) test
- North Star Ambulatory Assessment (NSAA) score
- Forced Vital Capacity (FVC) percent predicted

VI. Dosage/Administration

| Indication | Dose |
|-----------------------------|------------------------------------|
| Duchenne Muscular Dystrophy | 30 mg/kg intravenously once weekly |

VII. Billing Code/Availability Information

HCPCS Code:

• J1428 - Injection, eteplirsen, 10 mg; 1 billable unit = 10 mg

NDC:

- Exondys 51 100 mg/2 mL single-dose vial: 60923-0363-xx
- Exondys 51 500 mg/10 mL single-dose vial: 60923-0284-xx

VIII. References

- 1. Exondys 51 [package insert]. Cambridge, MA; Sarepta Therapeutics, Inc.; December 2024. Accessed April 2025.
- 2. Mendell JR, Rodino-Klapac LR, Sahenk Z, et al. Eteplirsen for the treatment of Duchenne muscular dystrophy. Ann Neurol. 2013 Nov;74(5):637-47. doi: 10.1002/ana.23982.
- 3. Mendell JR, Goemans N, Lowes LP, et al. Longitudinal effect of eteplirsen versus historical control on ambulation in Duchenne muscular dystrophy. Ann Neurol. 2016 Feb;79(2):257-71. doi: 10.1002/ana.24555.
- 4. Alfano LN, Charleston JS, Connolly AM, et al. Long-term treatment with eteplirsen in nonambulatory patients with Duchenne muscular dystrophy. Medicine (Baltimore). 2019 Jun;98(26): e15858. doi: 10.1097/MD.0000000000015858.

- 5. Topaloglu H, Gloss D, Moxley RT 3rd, et al. Practice guideline update summary: Corticosteroid treatment of Duchenne muscular dystrophy: Report of the Guideline Development Subcommittee of the American Academy of Neurology. Neurology. 2016 Jul 12;87(2):238.
- 6. Bushby K, Finkel R, Birnkrant DJ, et al. Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and pharmacological and psychosocial management. Lancet Neurol; 2010 Jan; 9(1):77-93.
- 7. Bushby K, Finkel R, Birnkrant DJ, et al. Diagnosis and management of Duchenne muscular dystrophy, part 2: implementation of multidisciplinary care. Lancet Neurol; 2010 Jan; 9(2):177-189.
- 8. Kinane TB, Mayer OH, Duda PW, et al. Long-Term Pulmonary Function in Duchenne Muscular Dystrophy: Comparison of Eteplirsen-Treated Patients to Natural History. Journal of Neuromuscular Diseases 5 (2018) 47–58.
- Institute for Clinical and Economic Review. Deflazacort, Eteplirsen, and Golodirsen for Duchenne Muscular Dystrophy: Effectiveness and Value. Final Evidence Report. August 15, 2019 https://icer-review.org/wp-content/uploads/2018/12/ICER_DMD-Final-Report_081519-1.pdf. Accessed December 2019.
- 10. Khan N, Eliopoulos H, et al on behalf of the Eteplirsen Investigators and the CINRG DNHS Investigators. Eteplirsen Treatment Attenuates Respiratory Decline in Ambulatory and Non-Ambulatory Patients with Duchenne Muscular Dystrophy. J. Neuromuscular Dis, vol. 6, no. 2, pp. 213-225, 2019.
- 11. Birnkrant DJ, Bushby K, Bann CM, et al. Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and neuromuscular, rehabilitation, endocrine, and gastrointestinal and nutritional management. Lancet Neurol 2018; 17:251.
- 12. Birnkrant DJ, Bushby K, Bann CM, et al. Diagnosis and management of Duchenne muscular dystrophy, part 2: respiratory, cardiac, bone health, and orthopaedic management. Lancet Neurol 2018; 17:347.
- 13. Moxley RT 3rd, Ashwal S, Pandya S, et al. Practice parameter: corticosteroid treatment of Duchenne dystrophy: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. Neurology. 2005;64:13–20.
- 14. Gloss D, Moxley RT 3rd, Ashwal S, Oskoui M. Practice guideline update summary: Corticosteroid treatment of Duchenne muscular dystrophy: Report of the Guideline Development Subcommittee of the American Academy of Neurology. Neurology. 2016 Feb 2;86(5):465-72. Doi: 10.1212/WNL.0000000000002337. Reaffirmed on January 22, 2022.



- 15. Darras BT, Urion DK, Ghosh PS. Dystrophinopathies. GeneReviews. www.ncbi.nlm.nih.gov/books/NBK1119/. Initial Posting: September 5, 2000; Last Revision: January 20, 2022. Accessed on July 1, 2024.
- 16. Childs, AM, Turner, C, Astin, R, et al. (2023). Development of respiratory care guidelines for Duchenne muscular dystrophy in the UK: key recommendations for clinical practice. thorax.
- 17. Landfeldt, Erik. Measuring health-related quality of life in Duchenne muscular dystrophy: Current perspectives and recommendations. Journal of the Neurological Sciences 446 (2023).

Appendix 1 - Covered Diagnosis Codes

| ICD-10 | ICD-10 Description |
|--------|---------------------------------------|
| G71.01 | Duchenne or Becker muscular dystrophy |

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

| Medicare Part B Administrative Contractor (MAC) Jurisdictions | | | | |
|---|-------------------------------------|---|--|--|
| Jurisdiction | Applicable State/US Territory | Contractor | | |
| E (1) | CA, HI, NV, AS, GU, CNMI | Noridian Healthcare Solutions, LLC | | |
| F (2 & 3) | AK, WA, OR, ID, ND, SD, MT, WY, UT, | Noridian Healthcare Solutions, LLC | | |
| 5 | KS, NE, IA, MO | Wisconsin Physicians Service Insurance Corp (WPS) | | |
| 6 | MN, WI, IL | National Government Services, Inc. (NGS) | | |
| H (4 & 7) | LA, AR, MS, TX, OK, CO, NM | Novitas Solutions, Inc. | | |
| 8 | MI, IN | Wisconsin Physicians Service Insurance Corp (WPS) | | |



| Medicare Part B Administrative Contractor (MAC) Jurisdictions | | | | |
|---|--|--|--|--|
| Jurisdiction | Applicable State/US Territory | Contractor | | |
| N (9) | FL, PR, VI | First Coast Service Options, Inc. | | |
| J (10) | TN, GA, AL | Palmetto GBA | | |
| M (11) | NC, SC, WV, VA (excluding below) | Palmetto GBA | | |
| L (12) | DE, MD, PA, NJ, DC (includes Arlington | Novitas Solutions, Inc. | | |
| | & Fairfax counties and the city of | | | |
| K (13 & 14) | NY, CT, MA, RI, VT, ME, NH | National Government Services, Inc. (NGS) | | |
| 15 | КҮ, ОН | CGS Administrators, LLC | | |

Policy Rationale: Exondys51 was reviewed by the Neighborhood Health Plan of Rhode Island Pharmacy & Therapeutics (P&T) Committee. Neighborhood adopted the following clinical coverage criteria to ensure that its members use Exondys51 according to Food and Drug Administration (FDA) approved labeling and/or relevant clinical literature. Neighborhood worked with network prescribers and pharmacists to draft these criteria. These criteria will help ensure its members are using this drug for a medically accepted indication, while minimizing the risk for adverse effects and ensuring more cost-effective options are used first, if applicable and appropriate. For Medicare members, these coverage criteria will only apply in the absence of National Coverage Determination (NCD) or Local Coverage Determination (LCD) criteria. Neighborhood will give individual consideration to each request it reviews based on the information submitted by the prescriber and other information available to the plan.