

Effective Date: 12/1/2023
Reviewed: 9/2023, 4/2024, 7/2025
Pharmacy Scope: Medicaid

JOENJA (leniolisib)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

Joenja is indicated for the treatment of activated phosphoinositide 3-kinase delta (PI3K δ) syndrome (APDS) in adults and pediatric patients 12 years of age and older.

All other indications are considered experimental/investigational and not medically necessary.

II. CRITERIA FOR INITIAL APPROVAL

Activated phosphoinositide 3-kinase delta (PI3K δ) syndrome (APDS)

Authorization of 6 months may be granted when all of the following criteria are met:

- A. Member is aged 12 to 75 years of age and weighing greater than or equal to 45 kg.
- B. This medication must be prescribed by or in consultation with an immunologist or a physician who specializes in the treatment of APDS.
- C. Member has documented APDS with a confirmed PIK3CD/PIK3R1 mutation without concurrent use of immunosuppressive medication.
- D. Documentation that the member has clinical manifestations compatible with APDS (e.g., nodal and/or extranodal lymphoproliferation, history of repeated oto-sino-pulmonary infections, autoimmunity [e.g., cytopenia], enteropathy, organ. dysfunction [e.g., lung, liver]) with or without the presence of ≥ 1 measurable nodal lesion on CT or MRI scan
- E. Documentation that the member has had an inadequate response, intolerance or contraindication to current standard of care for APDS (e.g., antimicrobial prophylaxis, immunoglobulin replacement therapy, immunosuppressive therapy).

III. CONTINUATION OF THERAPY

Authorization of 6 months may be granted for continued treatment in all members (including new members) who are currently receiving the requested medication and have documentation showing they are experiencing benefit from therapy as evidenced by disease stability or disease improvement. (i.e. reduced lymph node size, increased naïve B-cell percentage).

IV. QUANTITY LIMIT

Joenja has a quantity limit of 2 tablets/day.

V. REFERENCES

1. Joenja [package insert]. Warren, NJ: Pharming Technologies B.V.; March 2023.
2. Rao VK, Webster S, Šedivá A, et al. A randomized, placebo-controlled phase 3 trial of the PI3K δ inhibitor leniolisib for activated PI3K δ syndrome. *Blood*. 2023;141(9):971-983. doi:10.1182/blood.2022018546.