

Evolent Clinical Guideline 3101 for Hicon™ (sodium iodide I¹³¹)

Guideline Number: Evolent_CG_3101	<u>Applicable Codes</u>	
<p><i>"Evolent" refers to Evolent Health LLC and Evolent Specialty Services, Inc.</i></p> <p><i>© 2024 - 2025 Evolent. All rights Reserved.</i></p>		
Original Date: May 2024	Last Revised Date: October 2025	Implementation Date: October 2025

TABLE OF CONTENTS

STATEMENT	2
PURPOSE	2
INDICATIONS.....	2
THYROID CARCINOMA	2
CONTRAINDICATIONS/WARNINGS	2
EXCLUSION CRITERIA	3
CODING AND STANDARDS	3
CODES.....	3
APPLICABLE LINES OF BUSINESS	4
POLICY HISTORY.....	4
LEGAL AND COMPLIANCE.....	4
GUIDELINE APPROVAL	4
Committee.....	4
DISCLAIMER	4
REFERENCES.....	5

STATEMENT

Purpose

To define and describe the accepted indications for Hicon (sodium iodide I¹³¹) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

INDICATIONS

Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided

- The member has not experienced disease progression on the requested medication AND
- The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization AND
- Additional medication(s) are not being added to the continuation request.

Thyroid Carcinoma

- Hicon (sodium iodide I¹³¹) may be used in adult members in the following cases of thyroid carcinoma:
 - Member has a diagnosis of follicular, papillary, or oncocytic carcinoma: AND
 - Member had a thyroidectomy; OR
 - Member has locoregional, metastatic, or recurrent disease

CONTRAINDICATIONS/WARNINGS

- Contraindications
 - Patients with vomiting and diarrhea
 - Patients with thyroid malignancies shown to have no iodide uptake
 - Patients receiving concurrent anti-thyroid therapy
 - Pregnancy
 - Lactation

EXCLUSION CRITERIA

- Disease progression while taking Hicon (sodium iodide I¹³¹).
- Concurrent use with other anti-thyroid therapies.
- Dosing exceeds single dose limit of 100 millicuries (3,700 megabecquerels) for ablation of normal thyroid tissue, and 200 millicuries (7,400 megabecquerels) for subsequent ablation of metastases.
- Investigational use of Hicon (sodium iodide I¹³¹) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 - Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 - Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 - Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of < 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 - Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 - That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
 - That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
 - That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

CODING AND STANDARDS

Codes

- A9530 - Iodine i-131 sodium iodide solution, therapeutic, per millicurie

Applicable Lines of Business

<input type="checkbox"/>	CHIP (Children's Health Insurance Program)
<input checked="" type="checkbox"/>	Commercial
<input checked="" type="checkbox"/>	Exchange/Marketplace
<input checked="" type="checkbox"/>	Medicaid
<input type="checkbox"/>	Medicare Advantage

POLICY HISTORY

Date	Summary
October 2025	<ul style="list-style-type: none"> Corrected HCPC code
May 2025	<ul style="list-style-type: none"> Converted to new Evolent guideline template This guideline replaces UM ONC_1498 Hicon (sodium iodide I¹³¹) Updated references
May 2024	<ul style="list-style-type: none"> New policy

LEGAL AND COMPLIANCE

Guideline Approval

Committee

Reviewed / Approved by Evolent Specialty Clinical Guideline Review Committee

Disclaimer

Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment, and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. Evolent clinical guidelines contain guidance that requires prior authorization and service limitations. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as

required by applicable provider agreements and laws or regulations. Members should contact their Plan customer service representative for specific coverage information.

REFERENCES

1. Van Nostrand D. The benefits and risks of I-131 therapy in patients with well-differentiated thyroid cancer. *Thyroid*. 2009 Dec;19(12):1381-91. doi: 10.1089/thy.2009.1611.
2. Van Nostrand D. I131 ablation and treatment in well differentiated thyroid cancer. In: Van Nostrand D, Bloom G, Wartofsky L, Kulkarni KP, eds. *Thyroid Cancer: A Guide for Patients*. Pasadena, MD: Keystone Press; 2004;201-203.
3. Hicon prescribing information. Jubilant DraxImage Inc., dba Jubilant Radiopharm, Kirkland, Quebec H9H 4J4 Canada 2021.
4. Clinical Pharmacology Elsevier Gold Standard 2025.
5. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2025.
6. National Comprehensive Cancer Network. *Cancer Guidelines and Drugs and Biologics Compendium* 2025.
7. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs. Bethesda, MD 2025.
8. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol*. 2014 Apr 20;32(12):1277-80.
9. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
10. Current and Resolved Drug Shortages and Discontinuations Reported to the FDA: <http://www.accessdata.fda.gov/scripts/drugshortages/default.cfm>.