

Specialty Guideline Management

Mircera

Products Referenced by this Document

Drugs that are listed in the following table include both brand and generic and all dosage forms and strengths unless otherwise stated. Over-the-counter (OTC) products are not included unless otherwise stated.

Brand Name	Generic Name
Mircera	methoxy polyethylene glycol-epoetin beta

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications¹

Mircera is indicated for the treatment of anemia associated with chronic kidney disease (CKD) in:

- Adult patients on dialysis and adult patients not on dialysis.
- Pediatric patients 3 months to 17 years of age on dialysis or not on dialysis who are converting from another erythropoiesis-stimulating agent (ESA) after their hemoglobin level was stabilized with an ESA.

All other indications are considered experimental/investigational and not medically necessary.

Documentation

Submission of the following information is necessary to initiate the prior authorization review:

Continuation Requests

Reference number(s)
1618-A

Chart notes, medical records, or laboratory results of current (within the last 30 days) hemoglobin level.

Coverage Criteria

Note: Requirements regarding pretreatment hemoglobin level exclude values due to a recent transfusion. All members must be assessed for iron deficiency anemia and have adequate iron stores (defined as a serum transferrin saturation [TSAT] level greater than or equal to 20% within the prior 3 months) or are receiving iron therapy before starting Mircera. Members may not use Mircera concomitantly with other erythropoiesis stimulating agents (ESAs) or with hypoxia-inducible factor prolyl hydroxylase inhibitors (HIF-PHIs).

Anemia Due to Chronic Kidney Disease (CKD)^{1,2}

Authorization of 12 months may be granted for the treatment of anemia due to CKD in adult members with pretreatment hemoglobin (Hgb) less than 10 grams per deciliter (g/dL).

Authorization of 12 months may be granted for the treatment of anemia due to CKD in pediatric members 3 months to 17 years of age who are converting from another ESA after their hemoglobin level was stabilized (e.g., Hgb level of 10 to 12 g/dL) with an ESA.

Continuation Of Therapy

Note: Requirements regarding current hemoglobin level exclude values due to a recent transfusion. All members must be assessed for iron deficiency anemia and have adequate iron stores (defined as a serum transferrin saturation [TSAT] level greater than or equal to 20% with the prior 3 months) or are receiving iron therapy before continuation of treatment with Mircera. Members may not use Mircera concomitantly with other erythropoiesis-stimulating agents or with hypoxia-inducible factor prolyl hydroxylase inhibitors (HIF-PHIs).

Anemia Due to Chronic Kidney Disease (CKD)^{1,2}

Authorization of 12 months may be granted for continued treatment of anemia due to chronic kidney disease in members with current hemoglobin less than 12 g/dL.

References

1. Mircera [package insert]. St. Gallen, Switzerland: Vifor (International) Inc.; April 2024.
2. Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney Int.* 2012;Suppl 2:279-335.