

Cinryze® (C1 Esterase Inhibitor, Human) (Intravenous)

Effective Date: 01/01/2020

Review Date: 10/02/2019, 12/18/2019, 1/22/2020, 2/25/2021, 2/10/2022, 3/16/2023, 12/07/2023, 01/10/2024, 07/09/2025, 11/10/2025, 06/23/2026

Scope: Medicaid, Commercial, Medicare

I. Length of Authorization

Coverage will be provided for 6 months and may be renewed.

II. Dosing Limits

A. Max Units (per dose and over time) [HCPCS Unit]:

- 2,500 billable units per 30 days

III. Summary of Evidence

Cinryze is a plasma-derived replacement therapy indicated for routine prophylaxis against angioedema attacks in adults, adolescents, and pediatric patients (≥ 6 years old) with Hereditary Angioedema (HAE), a rare autosomal dominant disorder caused by C1 inhibitor deficiency. Cinryze's approval is based on a 24-week randomized, double-blind, placebo-controlled, crossover trial in 24 patients aged 9 to 73 years with a history of ≥ 2 HAE attacks per month. Patients receiving Cinryze (1,000 IU IV every 3–4 days) experienced a 52% reduction in attack frequency (mean 6.1 vs. 12.7; $p < 0.0001$), a 66% reduction in swelling days, shorter attack duration (2.1 vs. 3.4 days; $p = 0.0023$), and lower severity scores (1.3 vs. 1.9; $p = 0.0006$) compared to placebo. The most common adverse reactions ($\geq 5\%$) include headache, nausea, rash, vomiting, and fever. Serious adverse events such as thromboembolism have occurred, particularly in patients with known risk factors (e.g., indwelling catheters, obesity, oral contraceptives).

IV. Initial Approval Criteria¹

Coverage is provided in the following conditions:

Medicare members who have previously received this medication within the past 365 days are not subject to Step Therapy Requirements.

- Member is at least 6 years of age; **AND**

Universal Criteria ^{1,13}

- Must be prescribed by, or in consultation with, an allergist/immunologist or a physician who specializes in the management of HAE; **AND**
- Will not be used in combination with other prophylactic therapies targeting C1 inhibitor or kallikrein (i.e., Haegarda (C1 esterase inhibitor), Orladeyo (berotralstat), Takhzyro (lanadelumab-flyo), Andembry (garadacimab-gxii), Dawnzera (donidalorsen), etc); **AND**
- Other causes of angioedema have been ruled out (e.g., angiotensin-converting enzyme inhibitor [ACE-I] induced an angioedema, angioedema related to an estrogen containing drug, allergic angioedema).

Prophylaxis to prevent Hereditary Angioedema (HAE) attacks † Φ ¹⁻¹⁴

- Member has one of the clinical presentations listed below consistent with an HAE subtype§, which must be confirmed by repeat blood testing (treatment for acute attack should not be delayed for confirmatory testing); **AND**
 - Member is receiving treatment as short-term HAE prophylaxis prior to a procedure (i.e., dental or medical procedure); **OR**
 - Member requires long-term prophylactic treatment based on the provider's assessment of the patient's disease activity, quality of life, availability of health care resources, and/or failure to achieve adequate control by appropriate on-demand therapy [i.e., Ekterly (sebetralstat), Kalbitor(ecallantide), Icatibant, Ruconest (C1 esterase inhibitor) or Berinert (C1 esterase inhibitor), etc.]

HAE I (C1-Inhibitor deficiency) §

- Low C1 inhibitor (C1-INH) antigenic level (C1-INH antigenic level below the lower limit of normal as defined by the laboratory performing the test); **AND**
- Low C4 level (C4 below the lower limit of normal as defined by the laboratory performing the test); **AND**
- Low C1-INH functional level (C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test); **AND**
 - Member has a family history of HAE; **OR**
 - Acquired angioedema has been ruled out (i.e., member onset of symptoms occur prior to 30 years old, normal C1q levels, member does not have underlying disease such as lymphoma or benign monoclonal gammopathy [MGUS], etc.)

HAE II (C1-Inhibitor dysfunction) §

- Normal to elevated C1-INH antigenic level; **AND**
- Low C4 level (C4 below the lower limit of normal as defined by the laboratory performing the test); **AND**
- Low C1-INH functional level (C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test)

HAE with normal C1INH (formerly known as HAE III) §

- Prophylaxis for HAE with normal C1-INH is not routinely recommended and will be evaluated on a case-by-case basis
 - Prior to consideration of long-term prophylaxis, the member must have demonstrated:
 - An inadequate response or intolerance to an adequate trial of prophylactic therapy with an antifibrinolytic agent (e.g., tranexamic acid (TXA) or aminocaproic acid) and/or a 17 α -alkylated androgen (e.g., danazol) unless contraindicated. Female members may derive additional benefit from progestins^{15,16,17}; **AND**
 - Response to therapy from an agent indicated for the treatment of acute attacks (i.e., C1 esterase inhibitor, icatibant, ecallantide, etc.)

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ☐ Orphan Drug

V. Renewal Criteria¹

Coverage can be renewed based upon the following criteria:

- Member continues to meet the universal and other indication-specific relevant criteria identified in section IV; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: severe hypersensitivity reactions, serious thromboembolic events (arterial and venous), etc.; **AND**
 - Significant improvement in severity and duration of attacks have been achieved and sustained, **OR**
 - Member requires dose titration due to an inadequate response to therapy (> 1.0 HAE attack/month, regardless of severity/duration)

VI. Dosage/Administration¹

Indication	Dose
Prophylaxis of Hereditary Angioedema (HAE) attacks	Adult/adolescents (at least 12 years of age) 1,000 units by intravenous injection every 3 to 4 days – For members who have not responded adequately to initial dosing, doses up to 2,000 U (not exceeding 80 IU/kg) every 3 or 4 days may be considered based on individual member response. Pediatric members (6 to 11 years of age) 500 units by intravenous injection every 3 to 4 days – The dose may be adjusted according to individual member response, up to 1,000 U every 3 to 4 days. – One International Unit (IU) corresponds to the amount of C1 esterase inhibitor present in 1 mL of normal plasma as defined by the WHO international reference standard. Previously, the potency values were expressed in Units (U) and were relative to an in-house reference standard. A conversion factor of 0.83 can be used to recalculate the potency from U to IU, i.e., 100 U = 83 IU. – **Note: Patients may self-administer Cinryze after being instructed by their healthcare provider.

VII. Billing Code/Availability Information

HCPCS Code:

- J0598 – Injection, C1 esterase inhibitor (human), Cinryze, 10 units; 1 billable unit = 10 units

NDC:

- Cinryze 500 units single-dose vial: 42227-0081-xx

VIII. References

1. Cinryze [package insert]. Exton, PA; ViroPharma Biologics, Inc; November 2025. Accessed May 2026.
2. Lumry W, Manning ME, Hurewitz DS, et al, "Nanofiltered C1-Esterase Inhibitor for the Acute Management and Prevention of Hereditary Angioedema Attacks Due to C1-Inhibitor Deficiency in Children," J Pediatr, 2013, 162(5):1017-22.
3. Bowen T, Cicardi M, Farkas H, et al. Canadian 2003 International Consensus Algorithm For the Diagnosis, Therapy, and Management of Hereditary Angioedema. J Allergy Clin Immunol. 2004 Sep;114(3):629-37.
4. Bygum A, Andersen KE, Mikkelsen CS. Self-administration of intravenous C1-inhibitor therapy for hereditary angioedema and associated quality of life benefits. Eur J Dermatol. Mar-Apr 2009;19(2):147-151.
5. Bowen T, Cicardi M, Farkas H, et al. 2010 International consensus algorithm for the diagnosis, therapy and management of hereditary angioedema. Allergy Asthma Clin Immunol. 2010;6(1):24.
6. Craig T, Aygören-Pürsün E, Bork K, et al. WAO Guideline for the Management of Hereditary Angioedema. World Allergy Organ J. 2012 Dec;5(12):182-99.

7. Gompels MM, Lock RJ, Abinun M, et al. C1 inhibitor deficiency: consensus document. Clin Exp Immunol. 2005;139(3):379.
8. Betschel S, Badiou J, Binkley K, et al. Canadian hereditary angioedema guideline. Asthma Clin Immunol. 2014 Oct 24;10(1):50. doi: 10.1186/1710-1492-10-50.
9. Zuraw BL, Bernstein JA, Lang DM, et al. A focused parameter update: hereditary angioedema, acquired C1 inhibitor deficiency, and angiotensin-converting enzyme inhibitor-associated angioedema. J Allergy Clin Immunol. 2013 Jun;131(6):1491-3. doi: 10.1016/j.jaci.2013.03.034.
10. Zuraw BL, Banerji A, Bernstein JA, et al. US Hereditary Angioedema Association Medical Advisory Board 2013 recommendations for the management of hereditary angioedema due to C1 inhibitor deficiency. J Allergy Clin Immunol Pract. 2013 Sep-Oct;1(5):458-67.
11. Frank MM, Zuraw B, Banerji A, et al. Management of children with Hereditary Angioedema due to C1 Inhibitor deficiency. Pediatrics. 2016 Nov. 135(5)
12. Zuraw BL, Bork K, Binkley KE, et al. Hereditary angioedema with normal C1 inhibitor function: Consensus of an international expert panel. Allergy Asthma Proc. 2012;33 Suppl 1:145-156.
13. Maurer M, Mager M, Ansotegui I, et al. The international WAO/EAACI guideline for the management of hereditary angioedema-The 2017 revision and update. Allergy. 2018 Jan 10. doi: 10.1111/all.13384.
14. Lang DM, Aberer W, Bernstein JA, et al. International consensus on hereditary and acquired angioedema. Ann Allergy Asthma Immunol. 2012;109:395-402.
15. Wintemberger C, Boccon-Gibod I, Launay D, et al. Tranexamic acid as maintenance treatment for non-histaminergic angioedema: analysis of efficacy and safety in 37 patients. Clin Exp Immunol. 2014 Oct; 178(1): 112–117.
16. Saule C, Boccon-Gibod I, Fain O, et al. Benefits of progestin contraception in non-allergic angioedema. Clin Exp Allergy. 2013 Apr;43(4):475-82.
17. Frank MM, Sargent JS, Kane MA, et al. Epsilon aminocaproic acid therapy of hereditary angioneurotic edema; a double-blind study. N Engl J Med. 1972;286:808-812.
18. Zuraw B, Busse P, White M, et al. Efficacy and safety of long-term prophylaxis with C1 inhibitor (C1INH) concentrate in patients with hereditary angioedema (HAE). J Allergy Clin Immunol. 2008;121(2 Suppl 1):S272.
19. Aygören-Pürsün E, Soteres DF, Nieto-Martinez SA, et al. A randomized trial of human C1 inhibitor prophylaxis in children with hereditary angioedema. Pediatr Allergy Immunol. 2019;30(5):553-561.
20. Betschel S, Badiou J, Binkley K, et al. The International/Canadian Hereditary Angioedema Guideline. Allergy Asthma Clin Immunol. 2019; 15: 72. Published online 2019 Nov 25. doi: 10.1186/s13223-019-0376-8.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
D84.1	Defects in the complement system

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto Government Benefit Administrators, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

Policy Rationale: Cinryze was reviewed by the Neighborhood Health Plan of Rhode Island Pharmacy & Therapeutics (P&T) Committee. Neighborhood adopted the following clinical coverage criteria to ensure that its members use Cinryze according to Food and Drug Administration (FDA) approved labeling and/or relevant clinical literature. Neighborhood worked with network prescribers and pharmacists to draft these criteria. These criteria will help ensure its members are using this drug for a medically accepted indication, while minimizing the risk for adverse effects and ensuring more cost-effective options are used first, if applicable and appropriate. For Medicare members, these coverage criteria will only apply in the absence of National Coverage Determination (NCD) or Local Coverage Determination (LCD) criteria. Neighborhood will give individual consideration to each request it reviews based on the information submitted by the prescriber and other information available to the plan.