Evolent Clinical Guideline 3026 for Evdi/Yondelis™ (trabectedin)

Guideline Number:	Applicable Codes	
Evolent_CG_3026		
"Evolent" refers to Evolent Health LLC and Evolent Specialty Services, Inc. *© 2016 - 2026 Evolent. All rights Reserved.*		
Original Date:	**Last Revised Date:**	**Implementation Date:**
April 2016	July 2026	July 2026

TABLE OF CONTENTS

STATEMENT **2**
PURPOSE 2
INDICATIONS **2**
SOFT TISSUE SARCOMA 2
UTERINE SARCOMA 2
CONTRAINDICATIONS/WARNINGS **3**
EXCLUSION CRITERIA **3**
CODING AND STANDARDS **4**
CODES 4
APPLICABLE LINES OF BUSINESS 4
POLICY HISTORY **4**
LEGAL AND COMPLIANCE **4**
GUIDELINE APPROVAL 4
Committee *4*
DISCLAIMER 5
REFERENCES **6**

STATEMENT

Purpose

To define and describe the accepted indications for EvdiYondelis (trabectedin) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

INDICATIONS

Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided

- The member has not experienced disease progression on the requested medication AND
- The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization AND
- Additional medication(s) are not being added to the continuation request.

Soft Tissue Sarcoma

- The member has unresectable or metastatic soft tissue sarcoma (Leiomyosarcoma, liposarcoma, and translocation-related sarcomas) AND Evdi/Yondelis (trabectedin) may be used as monotherapy or in combination with doxorubicin as first line or subsequent therapy.
- NOTE: Evdi/Yondelis (trabectedin) is not supported by Evolent Policy for the treatment of other soft tissue sarcoma histologies that are not leiomyosarcoma, liposarcoma, and translocation-related sarcomas. This policy position is based on the T-SAR trial which demonstrated no clinically meaningful improvement in overall survival or progression free survival with Evdi/Yondelis (trabectedin) when compared to best supportive care for patients with advanced translocation-related sarcoma. Please refer to the alternative agents/regimens recommended by Evolent, including but not limited to regimens available at **Evolent Pathways**.

Uterine Sarcoma

- The member has unresectable or metastatic uterine leiomyosarcoma AND
- Evdi/Yondelis (trabectedin) is being used as single agent for members with disease

progression with prior anthracycline-based chemotherapy unless there is a contraindication/intolerance with prior anthracycline based therapy.

CONTRAINDICATIONS/WARNINGS

- Contraindications
  - Known, severe hypersensitivity (including anaphylaxis) to Evdi/Yondelis (trabectedin) or any component of the formulation.

EXCLUSION CRITERIA

- Disease progression while on Evdi/Yondelis (trabectedin).
- Dosing exceeds single dose limit of Evdi/Yondelis (trabectedin) 1.5 mg/m^2.
- Investigational use of Evdi/Yondelis (trabectedin) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
  - Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
  - Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
  - Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
  - Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
  - That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
  - That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
  - That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

CODING AND STANDARDS

Codes

- J9352 - Injection, trabectedin, 0.1 mg

Applicable Lines of Business

☐	CHIP (Children's Health Insurance Program)
☒	Commercial
☒	Exchange/Marketplace
☒	Medicaid
☐	Medicare Advantage

POLICY HISTORY

Date	Summary
July 2026	• Added new branded product "Evdi" to all relevant sections • Updated references
February 2026	• Updated references
February 2025	• Converted to new Evolent guideline template • This guideline replaces UM ONC_1290 Yondelis (trabectedin)
February 2024	• Updated NCH verbiage to Evolent

LEGAL AND COMPLIANCE

Guideline Approval

Committee

Reviewed / Approved by Evolent Specialty Services Clinical Guideline Review Committee

Disclaimer

Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment, and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. Evolent clinical guidelines contain guidance that requires prior authorization and service limitations. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. Members should contact their Plan customer service representative for specific coverage information.

REFERENCES

1. Pautier P, et al. French Sarcoma Group Clinical Trial, Doxorubicin alone versus doxorubicin with trabectedin followed by trabectedin alone as first-line therapy for metastatic or unresectable leiomyosarcoma (LMS-04): a randomised, multicentre, open-label phase 3 trial. *Lancet Oncol*. 2022 Aug;23(8):1044-1054. doi: 10.1016/S1470-2045(22)00380-1
2. Le Cesne A, et al. A randomized phase III trial comparing trabectedin to best supportive care in patients with pre-treated soft tissue sarcoma: T-SAR, a French Sarcoma Group trial. *Ann Oncol*. 2021 Aug;32(8):1034-1044. doi: 10.1016/j.annonc.2021.04.014
3. Patel S, et al. Overall survival and histology-specific subgroup analyses from a phase 3, randomized controlled study of trabectedin or dacarbazine in patients with advanced liposarcoma or leiomyosarcoma. *Cancer*. 2019 Aug 1;125(15):2610-2620. doi: 10.1002/cncr.32117
4. Demetri GD, et al. Efficacy and Safety of Trabectedin or Dacarbazine for Metastatic Liposarcoma or Leiomyosarcoma After Failure of Conventional Chemotherapy: Results of a Phase III Randomized Multicenter Clinical Trial. *J Clin Oncol*. 2016 Mar 10;34(8):786-93. doi: 10.1200/JCO.2015.62.4734
5. Yondelis prescribing information. Janssen Products, LP. Horsham, PA 2026.
6. Evdi prescribing information. Apotex Corp. Weston, FL 2026.
7. Clinical Pharmacology Elsevier Gold Standard 2026.
8. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2026.
9. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2026.
10. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs. Bethesda, MD 2026.
11. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol*. 2014 Apr 20;32(12):1277-80.
12. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf.