

SPECIALTY GUIDELINE MANAGEMENT

PCSK9i PRALUENT (alirocumab), REPATHA (evolocumab)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

- A. Members with established atherosclerotic cardiovascular disease.
- B. Members with an untreated LDL-C of greater than, or equal to, 190 mg/dL.

All other indications are considered experimental/investigational and not medically necessary.

II. CRITERIA FOR INITIAL APPROVAL

A. Clinical atherosclerotic cardiovascular disease (ASCVD)

Authorization of 12 months may be granted for treatment of ASCVD when all of the following criteria are met:

- 1. The member has a history of clinical atherosclerotic cardiovascular disease or has experienced a cardiovascular event
- 2. The member has a current LDL-C level greater than, or equal to, 70 mg/dL
- 3. The member is receiving maximally tolerated statin therapy or is statin intolerant

B. Primary or familial hyperlipidemia

Authorization of 12 months may be granted for treatment of primary or familial hyperlipidemia when all of the following criteria are met:

- 1. The member had an untreated (before any lipid lowering therapy) LDL-C level greater than, or equal to, 190 mg/dL
- 2. The member has a current LDL-C level greater than, or equal to, 100 mg/dL
- 3. The member is receiving maximally tolerated statin therapy or is statin intolerant

III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for members who are continuing therapy with a PCSK9i.

IV. REFERENCES

- 1. Repatha [package insert]. Thousand Oaks, CA: Amgen, Inc.; February 2021.
- 2. Praluent [package insert]. Bridgewater, NJ: sanofi-aventis U.S. LLC; September 2020.
- 3. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Circulation. 2014;129:S1-S45.

Reference number(s)
2574-A

4. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2017 Focused Update of the 2016 ACC expert consensus decision pathway on the role of non-statin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk: a report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol*. 2017;70:1785-822.
5. Cannon CP, Blazing MA, Giugliano RP, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med*. 2015;372:2387-97.
6. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 1 — full report. *J Clin Lipidol*. 2015;9:129–169.
7. Nordestgaard BG, Chapman MJ, Humphries SE, et al. Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: guidance for clinicians to prevent coronary heart disease. Consensus Statement of the European Atherosclerosis Society. *Eur Heart J*. 2013;34:3478–3490.
8. Cuchel M, Bruckert E, Ginsberg HN, et al. Homozygous familial hypercholesterolaemia: new insights and guidance for clinicians to improve detection and clinical management. A position paper from the Consensus Panel on Familial Hypercholesterolaemia of the European Atherosclerosis Society. *Eur Heart J*. 2014;35:2146-2157.
9. Raal FJ, Honarpour N, Blom DJ, et al. Inhibition of PCSK9 with evolocumab in homozygous familial hypercholesterolaemia (TESLA Part B): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2015;385:341-350.
10. Banach M, Rizzo M, Toth PP, et al. Statin intolerance – an attempt at a unified definition. Position paper from an International Lipid Expert Panel. *Arch Med Sci*. 2015;11:1-23.
11. Sabatine MC, Giugliano RP, Keech AC, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *New England Journal of Medicine*. 2017; Published online before print.
12. Hulten EA, Carbonaro S, Petrillo SP, et al. Prognostic value of cardiac computed tomography angiography. *Journal of the American College of Cardiology*. 2011;57(10):1237-1247.
13. Min JK, Labounty TM, Gomez MJ, et al. Incremental prognostic value of coronary computed tomographic angiography over coronary artery calcium score for risk predication of major adverse cardiac events in asymptomatic diabetic individuals. *Atherosclerosis*. 2014;232(2):298-304.
14. Robinson, J. G., Rogers, W. J., Nedergaard, B. S., Fialkow, J., Neutel, J. M., Ramstad, D., Somaratne, R., Legg, J. C., Nelson, P., Scott, R., Wasserman, S. M. and Weiss, R. (2014), Rationale and Design of LAPLACE-2: A Phase 3, Randomized, Double-Blind, Placebo- and Ezetimibe-Controlled Trial Evaluating the Efficacy and Safety of Evolocumab in Subjects With Hypercholesterolemia on Background Statin Therapy. *Clin Cardiol*, 37: 195–203.
15. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC guideline on the management of blood cholesterol: report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2018.