

Drug Policy:

Jemperli™ (dostarlimab-gxly)

POLICY NUMBER UM ONC_1433	SUBJECT Jemperli™ (dostarlimab-gxly)		DEPT/PROGRAM UM Dept	PAGE 1 OF 3
DATES COMMITTEE REVIEWED 06/09/21, 09/08/21, 11/15/21, 03/09/22	APPROVAL DATE March 9, 2022	EFFECTIVE DATE March 25, 2022	COMMITTEE APPROVAL DATES 06/09/21, 09/08/21, 11/15/21, 03/09/22	
PRIMARY BUSINESS OWNER: UM		COMMITTEE/BOARD APPROVAL Utilization Management Committee		
URAC STANDARDS HUM 1	NCQA STANDARDS UM 2		ADDITIONAL AREAS OF IMPACT	
CMS REQUIREMENTS	STATE/FEDERAL REQUIREMENTS		APPLICABLE LINES OF BUSINESS Commercial, Exchange, Medicaid	

I. PURPOSE

To define and describe the accepted indications for Jemperli (dostarlimab-gxly) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

New Century Health (NCH) is responsible for processing all medication requests from network ordering providers. Medications not authorized by NCH may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

II. INDICATIONS FOR USE/INCLUSION CRITERIA

A. PREFERRED MEDICATION GUIDANCE FOR INITIAL REQUEST:

1. When health plan Medicaid coverage provisions—including any applicable PDLs (Preferred Drug Lists)—conflict with the coverage provisions in this drug policy, health plan Medicaid coverage provisions take precedence per the [Preferred Drug Guidelines OR](#)
2. When health plan Exchange coverage provisions-including any applicable PDLs (Preferred Drug Lists)-conflict with the coverage provisions in this drug policy, health plan Exchange coverage provisions take precedence per the [Preferred Drug Guidelines OR](#)

3. For Health Plans that utilize NCH UM Oncology Clinical Policies as the initial clinical criteria, the [Preferred Drug Guidelines](#) shall follow [NCH L1 Pathways](#) when applicable, otherwise shall follow NCH drug policies **AND**
4. Continuation requests of previously approved, non-preferred medication are not subject to this provision **AND**
5. When applicable, generic alternatives are preferred over brand-name drugs.

B. Endometrial Carcinoma

1. Jemperli (dostarlimab-gxly) may be used as a single agent as subsequent line systemic therapy for unresectable or metastatic dMMR (deficient Mismatch Repair)/MSI-High (MicroSatellite Instability-High) endometrial carcinoma that has progressed following prior treatment with a platinum containing regimen **AND** the tumor is confirmed to be dMMR/MSI-High by any standard test.

C. Microsatellite Instability-High (MSI-H) or Mismatch Repair Deficient (dMMR) Solid Tumors

1. Jemperli (dostarlimab-gxly) may be used as monotherapy in members with recurrent, advanced, or metastatic solid tumors that have progressed following all satisfactory treatment alternatives and the solid tumor is positive for microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) as confirmed by any standard test.

III. EXCLUSION CRITERIA

- A. Disease progression while on or after Jemperli (dostarlimab-gxly) or on prior immunotherapy (anti-PD-L1 or PD-1 inhibitor).
- B. Concurrent use with other anti-cancer therapies (e.g., chemotherapy, endocrine therapy, targeted therapy, radiotherapy, or immunotherapy).
- C. Dosing exceeds single dose limit of Jemperli (dostarlimab-gxly) 500 mg every 3 weeks or 1,000 mg every 6 weeks.
- D. Investigational use of Jemperli (dostarlimab-gxly) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definition of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of < 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 4. Whether the experimental design, in light of the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.

6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

IV. MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

V. APPROVAL AUTHORITY

- A. Review – Utilization Management Department
- B. Final Approval – Utilization Management Committee

VI. ATTACHMENTS

- A. None

VII. REFERENCES

- A. Oaknin A, et al. GARNET trial: Clinical Activity and Safety of the Anti-Programmed Death 1 Monoclonal Antibody Dostarlimab for Patients With Recurrent or Advanced Mismatch Repair-Deficient Endometrial Cancer: A Nonrandomized Phase 1 Clinical Trial. *JAMA Oncol.* 2020 Nov 1;6(11):1766-1772.
- B. Jemperli PI prescribing information. GlaxoSmithKline LLC, Philadelphia, PA 2021.
- C. Clinical Pharmacology Elsevier Gold Standard 2022.
- D. Micromedex® Healthcare Series: Thomson Micromedex, Greenwood Village, CO 2022.
- E. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2022.
- F. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs Bethesda, MD 2022.
- G. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol.* 2014 Apr 20;32(12):1277-80.
- H. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.