

Drug Policy:

Abecma™ (idecabtagene vicleucel)

| | | | |
|---|--|---|---|
| POLICY NUMBER UM ONC_1429 | SUBJECT Abecma™ (idecabtagene vicleucel) | DEPT/PROGRAM UM Dept | PAGE 1 OF 4 |
| DATES COMMITTEE REVIEWED 05/12/21, 11/15/21, 02/09/22, 04/13/22, 05/11/22 | APPROVAL DATE May 11, 2022 | EFFECTIVE DATE May 27, 2022 | COMMITTEE APPROVAL DATES 05/12/21, 11/15/21, 02/09/22, 04/13/22, 05/11/22 |
| PRIMARY BUSINESS OWNER: UM | | COMMITTEE/BOARD APPROVAL Utilization Management Committee | |
| URAC STANDARDS HUM v8: UM 1-2; UM 2-1 | NCQA STANDARDS UM 2 | ADDITIONAL AREAS OF IMPACT | |
| CMS REQUIREMENTS | STATE/FEDERAL REQUIREMENTS | APPLICABLE LINES OF BUSINESS Commercial, Exchange, Medicaid | |

I. PURPOSE

To define and describe the accepted indications for Abecma (idecabtagene vicleucel) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

New Century Health (NCH) is responsible for processing all medication requests from network ordering providers. Medications not authorized by NCH may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

II. INDICATIONS FOR USE/INCLUSION CRITERIA

A. PREFERRED MEDICATION GUIDANCE FOR INITIAL REQUEST:

1. When health plan Medicaid coverage provisions—including any applicable PDLs (Preferred Drug Lists)—conflict with the coverage provisions in this drug policy, health plan Medicaid coverage provisions take precedence per the [Preferred Drug Guidelines OR](#)
2. When health plan Exchange coverage provisions—including any applicable PDLs (Preferred Drug Lists)—conflict with the coverage provisions in this drug policy, health plan Exchange coverage provisions take precedence per the [Preferred Drug Guidelines OR](#)
3. For Health Plans that utilize NCH UM Oncology Clinical Policies as the initial clinical criteria, the [Preferred Drug Guidelines shall follow NCH L1 Pathways](#)

(<http://pathways.newcenturyhealth.com/>) when applicable, otherwise shall follow NCH drug policies AND

4. Continuation requests of previously approved, non-preferred medication are not subject to this provision AND
5. When applicable, generic alternatives are preferred over brand-name drugs AND
6. When there is a documented drug shortage, disease progression, contraindication, or confirmed intolerance to a Preferred drug/regimen, per NCH Policy and Pathway, the available alternative product may be used if deemed medically appropriate and the indication is listed in a standard reference compendia or accepted peer review literature. For a list of current drug shortages, please refer to FDA drug shortage website in the reference section.

B. Multiple Myeloma

1. Abecma (idecabtagene vicleucel) may be used for adult members with relapsed/refractory multiple myeloma that have progressed on 4 or more lines of therapy AND
2. Members must have triple refractory disease as defined in the KaMMA trial: refractory to an immunomodulatory agent (e.g., lenalidomide, thalidomide, pomalidomide), a proteasome inhibitor (e.g., bortezomib, carfilzomib, ixazomib), and an anti-CD38 antibody (e.g., daratumumab, isatuximab).

III. EXCLUSION CRITERIA

- A. Disease progression while receiving Abecma (idecabtagene vicleucel).
- B. Concurrent use with other anti-myeloma therapy.
- C. The member does NOT have measurable disease defined as any of the following :
 1. Serum M-protein greater or equal to 1.0 g/dL
 2. Urine M-protein greater or equal to 200 mg/24 h
 3. Serum free light chain (FLC) assay: involved FLC level greater or equal to 10 mg/dL (100 mg/L) provided serum FLC ratio is abnormal.
- D. The member has solitary plasmacytomas or non-secretory myeloma without other evidence of measurable disease.
- E. The member does NOT have adequate bone marrow reserve defined by ALL of the following:
 1. Absolute neutrophil count (ANC) $\geq 1000/\mu\text{L}$
 2. Platelet Count $\geq 50,000/\mu\text{L}$.
- F. The member does NOT have adequate renal, hepatic, cardiac and pulmonary function defined as:
 1. Creatinine clearance $> 45 \text{ mL/min}$
 2. AST and/or ALT $\leq 2.5 \times \text{ULN}$ and total bilirubin $\leq 1.5 \times \text{ULN}$
 3. Cardiac ejection fraction $\geq 45\%$, no evidence of pericardial effusion as determined by an echocardiogram (ECHO), and no clinically significant pleural effusion.
- G. History or presence of CNS disorder.
- H. Prior allogeneic hematopoietic stem cell transplant (HSCT).
- I. Does not exceed duration limit as one time administration.
- J. Dosing exceeds single dose limit of Abecma (idecabtagene vicleucel) 460×10^6 CAR-positive T cells.

- K. Investigational use of Abecma (idecabtagene vicleucel) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definition of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of < 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 4. Whether the experimental design, in light of the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
 6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
 7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

IV. MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

V. APPROVAL AUTHORITY

- A. Review – Utilization Management Department
B. Final Approval – Utilization Management Committee

VI. ATTACHMENTS

- A. None

VII. REFERENCES

- A. Munshi NC, et al. KarMMa Trial. Idecabtagene Vicleucel in Relapsed and Refractory Multiple Myeloma. N Engl J Med. 2021 Feb 25;384(8):705-716.
- B. Raje N, et al. CRB-401 Trial. Anti-BCMA CAR T-Cell Therapy bb2121 in Relapsed or Refractory Multiple Myeloma. N Engl J Med. 2019 May 2;380(18):1726-1737.
- C. Abecma prescribing information. Celgene Corporation, Summit, NJ 2021.
- D. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol. 2014 Apr 20;32(12):1277-80.

- E. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services:
<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
- F. Current and Resolved Drug Shortages and Discontinuations Reported to the FDA:
<http://www.accessdata.fda.gov/scripts/drugshortages/default.cfm>.