

## SPECIALTY GUIDELINE MANAGEMENT

### CALQUENCE (acalabrutinib)

#### POLICY

##### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### A. FDA-Approved Indications

1. Mantle Cell Lymphoma  
Calquence is indicated for the treatment of adult patients with mantle cell lymphoma (MCL) who have received at least one prior therapy.
2. Chronic Lymphocytic Leukemia or Small Lymphocytic Lymphoma  
Calquence is indicated for the treatment of adult patients with chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL).

##### B. Compendial Use

1. Waldenström Macroglobulinemia/Lymphoplasmacytic Lymphoma
2. Gastric MALT Lymphoma/Non-gastric MALT Lymphoma
3. Nodal Marginal Zone Lymphoma
4. Splenic Marginal Zone Lymphoma

All other indications are considered experimental/investigational and not medically necessary.

##### II. CRITERIA FOR INITIAL APPROVAL

##### A. **Mantle cell lymphoma**

Authorization of 12 months may be granted for treatment of mantle cell lymphoma as a single agent when the member has received at least one prior therapy.

##### B. **Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL)**

Authorization of 12 months may be granted for treatment of CLL/SLL as a single agent or in combination with obinutuzumab.

##### C. **Waldenström Macroglobulinemia/Lymphoplasmacytic Lymphoma**

Authorization of 12 months may be granted for subsequent treatment of Waldenström Macroglobulinemia /Lymphoplasmacytic Lymphoma as a single agent.

##### D. **Gastric MALT Lymphoma/Non-gastric MALT Lymphoma/Nodal Marginal Zone Lymphoma/Splenic Marginal Zone Lymphoma**

Authorization of 12 months may be granted for treatment of gastric MALT lymphoma, non-gastric MALT lymphoma, nodal marginal zone lymphoma and splenic marginal zone lymphoma as subsequent therapy for members who are intolerant to or have contraindications to ibrutinib.

### III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization for an indication listed in Section II when there is no evidence of unacceptable toxicity or disease progression while on the current regimen.

### IV. REFERENCES

1. Calquence [package insert]. Wilmington, DE: AstraZeneca Pharmaceuticals LP; November 2019.
2. The NCCN Drugs & Biologics Compendium® © 2021 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed June 1, 2021.