# SPECIALTY GUIDELINE MANAGEMENT

## **CYSTADANE (betaine anhydrous)**

## POLICY

## I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

### A. FDA-Approved Indications

Cystadane is indicated for the treatment of homocystinuria to decrease elevated homocysteine blood concentrations in pediatric and adult patients. Included within the category of homocystinuria are:

- 1. Cystathionine beta-synthase (CBS) deficiency
- 2. 5,10-methylenetetrahydrofolate reductase (MTHFR) deficiency
- 3. Cobalamin cofactor metabolism (cbl) defect

#### B. Compendial Use

1. Methylmalonic acidemia with homocystinuria

All other indications are considered experimental/investigational and not medically necessary.

## **II. DOCUMENTATION**

Submission of the following information is necessary to initiate the prior authorization review:

- A. For cystathionine beta-synthase (CBS) deficiency, enzyme analysis of CBS activity or genetic testing results
- B. For 5,10-methylenetetrahydrofolate reductase (MTHFR) deficiency, enzyme analysis of MTHFR activity or genetic testing results
- C. For cobalamin cofactor metabolism (cbl) defect, genetic testing results

## **III. CRITERIA FOR INITIAL APPROVAL**

#### A. Homocystinuria

Authorization of 12 months may be granted for treatment of homocystinuria to decrease elevated homocysteine blood levels when all of the following criteria are met:

- 1. The member has one of the following types of homocystinuria and the diagnosis was confirmed by enzyme assay or genetic testing:
  - i. Cystathionine beta-synthase (CBS) deficiency
  - ii. 5,10-methylenetetrahydrofolate reductase (MTHFR) deficiency
  - iii. Cobalamin cofactor metabolism (cbl) defect
- 2. If the member has CBS deficiency, plasma methionine concentrations will be monitored and kept below 1,000 micromol/L through dietary modification, and if necessary, a reduction in Cystadane dose.

#### B. Methylmalonic acidemia with homocystinuria

#### Cystadane 2988-A SGM P2022

© 2022 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.



Authorization of 12 months may be granted for members who have a diagnosis of methylmalonic acidemia with homocystinuria.

#### **IV. CONTINUATION OF THERAPY**

#### A. Homocystinuria

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization for homocystinuria when both of the following criteria are met:

- 1. The total homocysteine level is undetectable or present only in small amounts, OR there is a substantial decrease in homocysteine levels and the dose will be increased until maximum tolerability or plasma total homocysteine is undetectable or present in only small amounts.
- 2. If the member has CBS deficiency, plasma methionine concentrations will be monitored and kept below 1,000 micromol/L through dietary modification, and if necessary, a reduction in Cystadane dose.

#### B. Methylmalonic acidemia with homocystinuria

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization for methylmalonic acidemia with homocystinuria who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

#### V. REFERENCES

- 1. Cystadane [package insert]. Lebanon, NJ: Recordati Rare Diseases, Inc.; October 2019.
- 2. Morris AA, Kožich V, Santra S, et al. Guidelines for the diagnosis and management of cystathionine betasynthase deficiency. *J Inherit Metab Dis.* 2017;40(1):49-74. doi:10.1007/s10545-016-9979-0.
- Sloan JL, Carrillo N, Adams D, et al. Disorders of Intracellular Cobalamin Metabolism. 2008 Feb 25 [Updated 2021 Dec 16]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2021.
- 4. Genetic and Rare Diseases Information Center. List of FDA Orphan Drugs. Methylmalonic Acidemia. https://rarediseases.info.nih.gov/diseases/fda-orphan-drugs/M. Accessed December 17, 2021.
- 5. National Organization for Rare Disorders (2003). *NORD guide to rare disorders*. Philadelphia: Lippincott Williams & Wilkins. Methylmalonic Acidemia. https://rarediseases.org/rare-diseases/acidemia-methylmalonic/. Accessed December 17, 2021.

Cystadane 2988-A SGM P2022

© 2022 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

