

## SPECIALTY GUIDELINE MANAGEMENT

### Cinacalcet

#### POLICY

##### 1. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### FDA-Approved Indication

Cinacalcet is indicated for secondary hyperparathyroidism (HPT) in adult patients with chronic kidney disease (CKD) on dialysis, hypercalcemia in adult patients with parathyroid carcinoma, and hypercalcemia in adult patients with primary HPT for whom parathyroidectomy would be indicated on the basis of serum calcium levels, but who are unable to undergo parathyroidectomy.

##### Compendial Use:

Tertiary hyperparathyroidism in post-kidney transplant patients not receiving dialysis

##### Limitations of use

Cinacalcet is not indicated for use in patients with CKD who are not on dialysis.

All other indications are considered experimental/investigational and are not a covered benefit.

##### 2. EXCLUSIONS:

Coverage will not be provided for members with a pretreatment serum calcium level (corrected for albumin) less than 8.4 mg/dL.

##### 3. CRITERIA FOR INITIAL APPROVAL

###### **A. Secondary hyperparathyroidism with CKD on dialysis**

Authorization of 12 months may be granted for members with secondary HPT with CKD on dialysis and a serum calcium level (corrected for albumin) greater than or equal to 8.4 mg/dL.

###### **B. Tertiary hyperparathyroidism in post-kidney transplant patients not receiving dialysis**

Authorization of 12 months may be granted for members with tertiary hyperparathyroidism not receiving dialysis who have had a kidney transplant and a serum calcium level (corrected for albumin) greater than or equal to 8.4 mg/dL.

Effective Date: 6/1/2020
Last Reviewed: 3/2020, 1/2021, 1/2022, 1/2023
Pharmacy Scope: Medicaid
Medical Scope: Medicaid, Commercial, MMP

### C. Parathyroid carcinoma

Authorization of 12 months may be granted for members with parathyroid carcinoma and a serum calcium level (corrected for albumin) greater than or equal to 8.4 mg/dL.

### D. Primary hyperparathyroidism

Authorization of 12 months may be granted for treatment of primary HPT when all of the following criteria are met:

1. The patient has a serum calcium level (corrected for albumin) greater than or equal to 8.4 mg/dL;
2. The patient is unable to undergo parathyroidectomy.

## 4. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for continuation of cinacalcet when either one of the two following criteria is met:

1. The patient has a serum calcium level (corrected for albumin) greater than or equal to 7.5 mg/dL
2. The patient has a serum calcium level (corrected for albumin) less than 7.5 mg/dL, and cinacalcet will be withheld until this level reaches 8 mg/dL

## 5. DOSAGE

Cinacalcet 30mg & 60mg have a quantity limit of 2 tablets per day

Cinacalcet 90mg has a quantity limit of 4 tablets per day

## 6. HCPCS codes

HCPCS Code	Description
J0604	Cinacalcet, oral, 1mg (for esrd on dialysis)

## REFERENCES

1. Sensipar [package insert]. Thousand Oaks, CA: Amgen, Inc.; December 2021.
2. Sterrett JR, Strom J, Stummvoll HK, Bahner U, Disney A, Soroka SD, et al. Cinacalcet HCl (Sensipar/Mimpara) is an effective chronic therapy for hemodialysis patients with secondary hyperparathyroidism. *Clin Nephrol*. 2007 Jul; 68: 10-7.
3. Kruse AE, Ensenberger U, Frey FJ, et al. The calcimimetic cinacalcet normalized serum calcium in renal transplant patients with persistent hyperparathyroidism. *Nephrol Dial Transplant*. 2005; 20: 1311-1314.