

<b>Policy Title:</b>	Triptodur (triptorelin) (Intramuscular)		
		<b>Department:</b>	PHA
<b>Effective Date:</b>	01/01/2020		
<b>Review Date:</b>	12/13/2019, 1/22/2020, 7/15/2021, 4/14/2022, 3/16/2023		
<b>Revision Date:</b>	12/13/2019, 1/22/2020, 7/15/2021		

**Purpose:** To support safe, effective and appropriate use of Triptodur (triptorelin).

**Scope:** Medicaid, Commercial, Medicare-Medicaid Plan (MMP)

**Policy Statement:**

Triptodur (triptorelin) is covered under the Medical Benefit when used within the following guidelines. Use outside of these guidelines may result in non-payment unless approved under an exception process.

**Procedure:**

Coverage of Triptodur (triptorelin) will be reviewed prospectively via the prior authorization process based on criteria below.

***Initial Criteria:***

**Central Precocious Puberty (CPP) † Φ:**

- Patient is between the ages of 2 and less than 13 years; AND
- Will not be used in combination with growth hormone; AND
- Onset of secondary sexual characteristics earlier than age 8 for girls and 9 for boys associated with pubertal pituitary gonadotropin activation; AND
- Diagnosis is confirmed by pubertal gonadal sex steroid levels and a pubertal luteinizing hormone (LH) response to stimulation by native GnRH; AND
- Bone age advanced greater than 2 standard deviations (SD) beyond chronological age; AND
- Tumor has been ruled out by lab tests such as diagnostic imaging of the brain (to rule out intracranial tumor), pelvic/testicular/adrenal ultrasound (to rule out steroid secreting tumors), and human chorionic gonadotropin levels (to rule out a chorionic gonadotropin secreting tumor); AND
- Patient must have a documented failure, intolerance or contraindication to Trelstar (triptorelin pamoate)
- MMP members who have previously received this medication within the past 365 days are not subject to Step Therapy Requirements

### **Gender Dysphoria (formerly Gender Identity Disorder) ‡**

- Patient has a diagnosis of gender dysphoria as confirmed by a qualified mental health professional (MHP)\*\* OR the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) Criteria §; **AND**
- A qualified MHP\*\* has confirmed all of the following:
  - Patient has demonstrated a long-lasting and intense pattern of gender nonconformity or gender dysphoria (whether suppressed or expressed); **AND**
  - Gender dysphoria worsened with the onset of puberty; **AND**
  - Any coexisting psychological, medical, or social problems that could interfere with treatment (e.g., that may compromise treatment adherence) have been addressed, such that the adolescent's situation and functioning are stable enough to start treatment; **AND**
  - Patient has sufficient mental capacity to give informed consent to this (reversible) treatment; **AND**
- Patient must have a documented failure, intolerance, or contraindication to Lupron Depot (Leuprolide acetate)
- Patient has been informed of the effects and side effects of treatment (including potential loss of fertility if the individual subsequently continues with sex hormone treatment) and options to preserve fertility; **AND**
- Patient has given informed consent and (particularly when the adolescent has not reached the age of legal medical consent, depending on applicable legislation) the parents or other caretakers or guardians have consented to the treatment and are involved in supporting the adolescent throughout the treatment process; **AND**
- A pediatric endocrinologist or other clinician experienced in pubertal assessment has confirmed all the following:
  - Agreement in the indication for treatment; **AND**
  - Puberty has started in the adolescent (e.g., Tanner stage  $\geq$ G2/B2); **AND**
  - There are no medical contraindications to treatment

#### **\*\* Definition of a qualified mental health professional**

- A master's degree or its equivalent in a clinical behavioral science field. This degree or a more advanced one should be granted by an institution accredited by the appropriate national or regional accrediting board. The mental health professional should also have documented credentials from the relevant licensing board or equivalent; **AND**
- Competence in using the Diagnostic Statistical Manual of Mental Disorders and/or the International Classification of Diseases for diagnostic purposes; **AND**
- Ability to recognize and diagnose co-existing mental health concerns and to distinguish these from gender dysphoria; **AND**
- Knowledgeable about gender nonconforming identities and expressions, and the assessment and treatment of gender dysphoria; **AND**
- Continuing education in the assessment and treatment of gender dysphoria. This may include attending relevant professional meetings, workshops, or seminars; obtaining supervision from a mental health professional with relevant experience; or participating in research related to gender nonconformity and gender dysphoria.

#### § DSM-V Criteria for Gender Dysphoria

- A marked incongruence between one's experienced/expressed gender and natal gender of at least 6mo in duration, as manifested by at least TWO of the following:
  - A marked incongruence between one's experienced/expressed gender and primary and/or secondary sex characteristics (or in young adolescents, the anticipated secondary sex characteristics)
  - A strong desire to be rid of one's primary and/or secondary sex characteristics because of a marked incongruence with one's experienced/expressed gender (or in young adolescents, a desire to prevent the development of the anticipated secondary sex characteristics)
  - A strong desire for the primary and/or secondary sex characteristics of the other gender
  - A strong desire to be of the other gender (or some alternative gender different from one's designated gender)
  - A strong desire to be treated as the other gender (or some alternative gender different from one's designated gender)
  - A strong conviction that one has the typical feelings and reactions of the other gender (or some alternative gender different from one's designated gender); **AND**
- The condition is associated with clinically significant distress or impairment in social, occupational, or other important areas of functioning; **AND**
- Specify one of the following:
  - The condition exists with a disorder of sex development, **OR**
  - The condition is post-transitional, in that the individual has transitioned to full-time living in the desired gender (with or without legalization of gender change) and has undergone (or is preparing to have) at least one sex-related medical procedure or treatment regimen—namely, regular sex hormone treatment or gender reassignment surgery confirming the desired gender (e.g., penectomy, vaginoplasty in natal males; mastectomy or phalloplasty in natal females).

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ☐ Orphan Drug

#### ***Continuation of Therapy Criteria:***

- Patient continues to meet initial criteria; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include psychiatric events (e.g., emotional lability including crying, irritability, impatience, anger, and aggression), convulsions, etc.

#### **Central Precocious Puberty (CPP)**

- Patient is less than 13 years of age; **AND**
- Disease response as indicated by lack of progression or stabilization of secondary sexual characteristics, decrease in height velocity, a decrease in the ratio of bone age to chronological age (BA:CA), and improvement in final height prediction

#### **Gender Dysphoria**

- Patient has shown a beneficial response to treatment as evidenced by routine monitoring of clinical pubertal development and applicable laboratory parameters

#### **Coverage durations:**

- Initial coverage: 6 months
- Continuation of therapy coverage: 6 months

\*\*\* Requests will also be reviewed to National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) if applicable.\*\*\*

**Dosage/Administration:**

Indication	Dose	Maximum dose (1 billable unit = 3.75 mg)
CPP and Gender Dysphoria	22.5 mg administered by a healthcare professional as a single intramuscular injection once every 24 weeks	6 billable units per 168 days

**Investigational use:** All therapies are considered investigational when used at a dose or for a condition other than those that are recognized as medically accepted indications as defined in any one of the following standard reference compendia: American Hospital Formulary Service Drug information (AHFS-DI), Thomson Micromedex DrugDex, Clinical Pharmacology, Wolters Kluwer Lexi-Drugs, or Peer-reviewed published medical literature indicating that sufficient evidence exists to support use. Neighborhood does not provide coverage for drugs when used for investigational purposes.

**Applicable Codes:**

Below is a list of billing codes applicable for covered treatment options. The below tables are provided for reference purposes and may not be all-inclusive. Requests received with codes from tables below do not guarantee coverage. Requests must meet all criteria provided in the procedure section.

The following HCPCS/CPT codes are:

HCPCS/CPT Code	Description
J3316	Injection Triptorelin, extended release

**References:**

1. Triptodur [package insert]. Atlanta, GA; Arbor Pharmaceutical, LLC; December 2022. Accessed March 2023.
2. Klein K, et al. Efficacy and safety of triptorelin 6-month formulation in patients with central precocious puberty. J Pediatr Endocrinol Metab. 2016;29(11):1241-1248.
3. Carel JC, Eugster EA, Rogol A, et al. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. Pediatrics. 2009; 123(4):e752.
4. Brito VN, Spinola-Castro AM, Kochi C, et al. Central precocious puberty: revisiting the diagnosis and therapeutic management. Arch Endocrinol Metab. 2016 Apr;60(2):163-72
5. First Coast Service Options, Inc. Local Coverage Determination (LCD): Luteinizing Hormone-Releasing Hormone (LHRH) Analogs (L33685). Centers for Medicare & Medicaid Services, Inc. Updated on 5/7/2018 with effective date 3/15/2018. Accessed July 2018.
6. Novitas Solutions, Inc. Local Coverage Determination (LCD): Luteinizing Hormone-Releasing Hormone (LHRH) Analogs (L34822). Centers for Medicare & Medicaid Services, Inc. Updated on 12/09/2014 with effective date 10/01/2015. Accessed July 2018.
7. Carel JC, Blumberg J, Seymour C, et al. Three-month sustained-release triptorelin (11.25 mg) in the treatment of central precocious puberty. Eur J Endocrinol 2006; 154:119.

8. Klein K, Yang J, Aisenberg J, et al. Efficacy and safety of triptorelin 6-month formulation in patients with central precocious puberty. *J Pediatr Endocrinol Metab* 2016; 29:1241.
9. Bertelloni S, Massart F, Einaudi S, et al. Central Precocious Puberty: Adult Height in Girls Treated with Quarterly or Monthly Gonadotropin-Releasing Hormone Analog Triptorelin. *Horm Res Paediatr* 2015; 84:396.
10. Krishna KB, Fuqua JS, Rogol AD, et al. Use of Gonadotropin-Releasing Hormone Analogs in Children: Update by an International Consortium *Horm Res Paediatr* 2019;91:357–372
11. Kaplowitz Bloch C; Section on Endocrinology, American Academy of Pediatrics. Evaluation and Referral of Children With Signs of Early Puberty. *Pediatrics*. 2016 Jan;137(1). Doi: 10.1542/peds.2015-3732. Epub 2015 Dec 14.
12. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2017; 102:3869.
13. The World Professional Association for Transgender Health (WPATH), Standards of Care for the Health of Transsexual, and Gender Nonconforming People. Seventh Version. July 2012. Available at: [https://www.wpath.org/media/cms/Documents/SOC%20v7/SOC%20V7\\_English2012.pdf? t=1613669341](https://www.wpath.org/media/cms/Documents/SOC%20v7/SOC%20V7_English2012.pdf? t=1613669341)
14. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5<sup>th</sup> ed. Arlington, VA: American Psychiatric Association Publishing.
15. Schagen SE, Cohen-Kettenis PT, Delemarre-van de Waal HA, et al: Efficacy and safety of gonadotropin-releasing hormone agonist treatment to suppress puberty in gender dysphoric adolescents. *J Sex Med* 2016; 13(7):1125-1132.