



Hemophilia Products – Anti-Inhibitor Antibody: Hemlibra (emicizumab-kxwh)

(Subcutaneous)

Effective date: 01/01/2020

Review date: 12/18/19, 1/22/20, 2/25/2021, 06/24/2021, 6/16/2022, 6/22/2023

Scope: Medicaid*, Commercial*, Medicare-Medicaid Plan (MMP)

*(Medication only available on the Pharmacy Benefit)

I. Length of Authorization

Unless otherwise specified*, the initial authorization will be provided for 3 months and may be renewed every 12 months thereafter.

Note: The cumulative amount of medication the patient has on-hand will be taken into account for authorizations.

II. Dosing Limits

A. Quantity Limit (max daily dose) [Pharmacy Benefit]:

Loading Dose:

345mg weekly x 4 doses

Maintenance Dose:

- 1.5 mg/kg weekly = 180 mg weekly
- 3mg/kg every 2 weeks = 345mg every 2 weeks
- 6mg/kg every 4 weeks = 690mg every 4 weeks

B. Max Units (per dose and over time) [Medical Benefit]:

Loading Dose:


• 690 billable units (BU) weekly x 4 doses

Maintenance Dose:

- 1.5 mg/kg weekly = 360 BU weekly
- 3mg/kg every 2 weeks = 690 BU every 2 weeks
- 6mg/kg every 4 weeks = 1380 BU every 4 weeks

^{*} Initial and renewal authorization periods may vary by specific covered indication

^{***} Requests will also be reviewed to National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) if applicable.***





<u>Note</u>: Patient must be dosed at a frequency that will produce the least wastage per dose based on available vial sizes of 30 mg, 60 mg, 105 mg, 150 mg, and 300mg.

III. Initial Approval Criteria

Hemophilia A (congenital factor VIII deficiency) with inhibitors \dagger Φ

- Diagnosis of congenital factor VIII deficiency has been confirmed by blood coagulation testing; AND
- Confirmation the patient has inhibitors to Factor VIII; AND
- Used as routine prophylaxis to prevent or reduce the frequency of bleeding episodes; AND
- Not used in combination with Immune Tolerance Induction (ITI); AND
 - o Patient has at least two documented episodes of spontaneous bleeding into joints; **OR**
 - o Patient has documented trial and failure of Immune Tolerance Induction (ITI); **OR**
 - o Patient has documented trial and failure of or is currently on routine prophylaxis with a bypassing agent (i.e., NovoSeven, FEIBA).

Hemophilia A (congenital factor VIII deficiency) without inhibitors † Φ

- Diagnosis of congenital factor VIII deficiency has been confirmed by blood coagulation testing; AND
- Used for routine prophylaxis: **AND**
 - o Patient must have severe hemophilia A (factor VIII level of <1%); **OR**
 - Patient has at least two documented episodes of spontaneous bleeding into joints; AND
- Patient is not a suitable candidate for treatment with shorter half-life Factor VIII (recombinant) products at a
 total weekly dose of 100 IU/kg or less (as attested by the prescribing physician with appropriate clinical
 rationale)
- † FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Orphan Drug

IV. Dispensing Requirements for Rendering Providers (Hemophilia Management Program)

- Prescriptions cannot be filled without an expressed need from the patient, caregiver or prescribing practitioner. Auto-filling is not allowed.
- Monthly, rendering provider must submit for authorization of dispensing quantity before delivering factor product.
- The cumulative amount of medication(s) the patient has on-hand should be taken into account when dispensing factor product.
- Dispensing requirements for renderings providers are a part of the hemophilia management program. This
 information is not meant to replace clinical decision making when initiating or modifying medication therapy
 and should only be used as a guide.



V. Renewal Criteria

Coverage can be renewed based upon the following criteria:

- Patient continues to meet criteria identified in section III; AND
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: thrombotic
 microangiopathy and thrombotic events, thromboembolic events (thromboembolism, pulmonary embolism),
 development of neutralizing antibodies (inhibitors), etc.; AND
- Any increases in dose must be supported by an acceptable clinical rationale (i.e. weight gain, half-life study results, increase in breakthrough bleeding when patient is fully adherent to therapy, etc.); **AND**
- Patient has demonstrated a beneficial response to therapy (i.e., the frequency of bleeding episodes has
 decreased from pre-treatment baseline); AND
- The cumulative amount of medication(s) the patient has on-hand will be taken into account when authorizing.

Routine prophylaxis to prevent or reduce the frequency of bleeding episode

• Renewals will be approved for a 12 month authorization period

Dosage/Administration

Indication	Dose
Routine Prophylaxis in	3 mg/kg by subcutaneous injection once weekly for the
Congenital Hemophilia	first 4 weeks, followed by 1.5 mg/kg once weekly, 3 mg/kg every two weeks, or 6 mg/kg
A with or without	every four weeks
inhibitors	

VI. Billing Code/Availability Information

HCPCS Code:

• J7170 - Injection, emicizumab-kxwh, 0.5 mg; 1 billable unit = 0.5 mg

NDC:

Drug	Strength	Form	NDC
	30 mg/mL	SDV	50242-0920-xx
	60 mg/0.4 mL	SDV	50242-0921- xx
Hemlibra	105 mg/0.7 mL	SDV	50242-0922- xx
	150 mg/mL	SDV	50242-0923- xx
	300 mg/2 mL	SDV	50242-0930-xx

VII. References

1. Hemlibra [package insert]. South San Francisco, CA; Genentech, Inc. March 2023. Accessed May 2023.



- 2. MASAC RECOMMENDATIONS CONCERNING PRODUCTS LICENSED FOR THE TREATMENT OF HEMOPHILIA AND OTHER BLEEDING DISORDERS. 2016 National Hemophilia Foundation. MASAC Document #249; October 2016. Available at: http://www.hemophilia.org. Accessed January 2018.
- 3. Guidelines for the Management of Hemophilia. 2nd Edition. World Federation of Hemophilia. 2013. Available at: https://www1.wfh.org/publication/files/pdf-1472.pdf. Accessed January 2019.
- 4. Annual Review of Factor Replacement Products. Oklahoma Health Care Authority Review Board. Updated April 2016. Access June 2016.
- 5. Graham A1, Jaworski K. Pharmacokinetic analysis of anti-hemophilic factor in the obese patient. Haemophilia. 2014 Mar;20(2):226-9.
- 6. Croteau SE1, Neufeld EJ. Transition considerations for extended half-life factor products. Haemophilia. 2015 May;21(3):285-8.
- 7. Mingot-Castellano, et al. Application of Pharmacokinetics Programs in Optimization of Haemostatic Treatment in Severe Hemophilia a Patients: Changes in Consumption, Clinical Outcomes and Quality of Life. Blood. 2014 December; 124 (21).
- 8. MASAC RECOMMENDATION CONCERNING PROPHYLAXIS. 2016 National Hemophilia Foundation. MASAC Document #241; February 2016. Available at: http://www.hemophilia.org. Accessed January 2019.
- UKHCDO protocol for first line immune tolerance induction for children with severe haemophilia A: A
 protocol from the UKHCDO Inhibitor and Paediatric Working Parties. 2017. Available at:
 http://www.ukhcdo.org/guidelines. Accessed January 2019.

Appendix 1 – Covered Diagnosis Codes

Hemlibra

ICD-10	ICD-10 Description
D66	Hereditary factor VIII deficiency



Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications may be covered at the discretion of the health plan.


Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

Jurisdiction(s): N	NCD/LCD Document (s): A56482	
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56482&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP		
Jurisdiction(s): J,M	NCD/LCD Document (s): A56065	
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56065&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP		

	Jurisdiction(s): H,L	NCD/LCD Document (s): A56433	
1	https://www.cms.gov/medicare-coverage-database/new-search/search-		
1	results.aspx?keyword=a56433&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2		
9	C6%2C3%2C5%2C1%2CF%2	<u>ecp</u>	

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD): N/A

	Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor	
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC	
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC	
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)	
6	MN, WI, IL	National Government Services, Inc. (NGS)	
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.	
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)	
N (9)	FL, PR, VI	First Coast Service Options, Inc.	
J (10)	TN, GA, AL	Palmetto GBA, LLC	
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC	





	Medicare Part B Administrative Contractor (MAC) Jurisdictions			
Jurisdiction	Applicable State/US Territory	Contractor		
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	КҮ, ОН	CGS Administrators, LLC		