

Reference number
2017-A

SPECIALTY GUIDELINE MANAGEMENT

COSENTYX (secukinumab)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

1. Moderate to severe plaque psoriasis in adult patients who are candidates for systemic therapy or phototherapy
2. Adults with active psoriatic arthritis (PsA)
3. Adults with active ankylosing spondylitis
4. Adults with active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation

All other indications are considered experimental/investigational and not medically necessary.

II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

- A. Plaque psoriasis
 1. Initial requests:
 - i. Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected.
 - ii. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 2. Continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.
- B. Psoriatic arthritis: For continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- C. Ankylosing spondylitis and axial spondyloarthritis:
 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy if not advisable, documentation of clinical reason to avoid therapy.
 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.

III. CRITERIA FOR INITIAL APPROVAL

Reference number
2017-A

A. Moderate to severe plaque psoriasis (PsO)

1. Authorization of 12 months may be granted for members who have previously received Otezla or a biologic indicated for the treatment of moderate to severe plaque psoriasis.
2. Authorization of 12 months may be granted for treatment of moderate to severe plaque psoriasis in members when any of the following criteria is met:
 - i. Crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - ii. At least 10% of the body surface area (BSA) is affected
 - iii. At least 3% of body surface area (BSA) is affected and the member meets any of the following criteria:
 - a. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine or acitretin.
 - b. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine and acitretin (see Appendix).

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for treatment of active psoriatic arthritis.

C. Active ankylosing spondylitis (AS) and active axial spondyloarthritis

1. Authorization of 12 months may be granted for members who have previously received a biologic indicated for active ankylosing spondylitis or active axial spondyloarthritis.
2. Authorization of 12 months may be granted for treatment of active ankylosing spondylitis or active axial spondyloarthritis when any of the following criteria is met:
 - i. Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
 - ii. Member has an intolerance or contraindication to two or more NSAIDs.

IV. CONTINUATION OF THERAPY

A. Moderate to severe plaque psoriasis (PsO)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

1. Reduction in body surface area (BSA) affected from baseline
2. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active psoriatic arthritis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Skin and/or nail involvement

Reference number
2017-A

C. Active ankylosing spondylitis (AS) and active axial spondyloarthritis

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active ankylosing spondylitis or active axial spondyloarthritis and who achieve or maintain positive clinical response with the requested medication as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Functional status
2. Total spinal pain
3. Inflammation (e.g., morning stiffness)

V. OTHER

For all indications: Member has had a documented negative TB test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for persons who are naïve to biologic DMARDs or targeted synthetic DMARDs associated with an increased risk of TB, and repeated yearly for members with risk factors** for TB that are continuing therapy with biologics.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

** Risk factors for TB include: Persons with close contact to people with infectious TB disease; persons who have recently immigrated from areas of the world with high rates of TB (e.g., Africa, Asia, Eastern Europe, Latin America, Russia); children less than 5 years of age who have a positive TB test; groups with high rates of TB transmission (e.g., homeless persons, injection drug users, persons with HIV infection); persons who work or reside with people who are at an increased risk for active TB (e.g., hospitals, long-term care facilities, correctional facilities, homeless shelters).

For all indications: Member cannot use the requested medication concomitantly with any other biologic DMARD or targeted synthetic DMARD.

VI. DOSAGE AND ADMINISTRATION

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines

VII. APPENDIX

Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine or Acitretin

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Drug interaction
4. Cannot be used due to risk of treatment-related toxicity
5. Pregnancy or currently planning pregnancy
6. Significant comorbidity prohibits use of systemic agents (examples include liver or kidney disease, blood dyscrasias, uncontrolled hypertension)

Reference number
2017-A

VIII. REFERENCES

1. Cosentyx [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; June 2020.
2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.
3. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis*. 2020;79(6):700-712.
4. McInnes IB, Mease PJ, Kirkham B, et al. Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (FUTURE 2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2015;386(9999):1137-46.
5. Braun J, van den Berg R, Baraliakos, X et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis*. 2011;70:896–904.
6. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Rheumatol*. 2019;71(10):1285-1299.
7. Baeten D, Sieper J, Braun J, et al. Secukinumab, an Interleukin-17A Inhibitor, in Ankylosing Spondylitis. *N Engl J Med*. 2015;373(26):2534-48.
8. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80(4):1029-1072.
9. Tuberculosis (TB). TB risk factors. Centers for Disease Control and Prevention. Retrieved on 14 January 2021 from: <https://www.cdc.gov/tb/topic/basics/risk.htm>.
10. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheum*. 2018;71:5-32.