

Reference number(s)
1651-A

SPECIALTY GUIDELINE MANAGEMENT

REVATIO (sildenafil) LIQREV (sildenafil) sildenafil

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Revatio/Liqrev/sildenafil is indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1) in adults to improve exercise ability and delay clinical worsening.
2. Revatio/sildenafil is indicated in pediatric patients 1 to 17 years old for the treatment of pulmonary arterial hypertension (PAH) (WHO Group I) to improve exercise ability and, in pediatric patients too young to perform standardized exercise testing, pulmonary hemodynamics thought to underly improvements in exercise.

B. Compendial Uses

1. Secondary Raynaud's phenomenon
2. Pulmonary arterial hypertension (PAH) (WHO Group I) in pediatric members less than 1 year of age

All other indications are considered experimental/investigational and not medically necessary.

II. PRESCRIBER SPECIALTIES

This medication must be prescribed by or in consultation with a pulmonologist or cardiologist for the diagnosis of pulmonary arterial hypertension (PAH).

III. CRITERIA FOR INITIAL APPROVAL

A. **Pulmonary Arterial Hypertension (PAH)**

Authorization of 12 months may be granted for treatment of PAH when ALL of the following criteria are met:

1. Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
2. PAH was confirmed by either criterion (i) or criterion (ii) below:
 - i. Pretreatment right heart catheterization with all of the following results:
 - a. Mean pulmonary arterial pressure (mPAP) > 20 mmHg
 - b. Pulmonary capillary wedge pressure (PCWP) ≤ 15 mmHg
 - c. Pulmonary vascular resistance (PVR) ≥ 3 Wood units in adult patients or pulmonary vascular resistance index (PVRI) ≥ 3 Wood units x m² in pediatric patients

Reference number(s)
1651-A

- ii. For infants less than one year of age, PAH was confirmed by Doppler echocardiogram if right heart catheterization cannot be performed.

B. Secondary Raynaud’s Phenomenon

Authorization of 12 months may be granted for treatment of secondary Raynaud’s phenomenon when the member has had an inadequate response to one of the following medications:

- 1. Calcium channel blockers
- 2. Angiotensin II receptor blockers
- 3. Selective serotonin reuptake inhibitors
- 4. Alpha blockers
- 5. Topical nitrates

IV. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for members with an indication listed in Section III who are currently receiving the requested medication through a paid pharmacy or medical benefit, and who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

V. APPENDIX

WHO Classification of Pulmonary Hypertension (PH)

1 PAH

- 1.1 Idiopathic (PAH)
- 1.2 Heritable PAH
- 1.3 Drug- and toxin-induced PAH
- 1.4. PAH associated with:
 - 1.4.1 Connective tissue diseases
 - 1.4.2 Human Immunodeficiency Virus (HIV) infection
 - 1.4.3 Portal hypertension
 - 1.4.4 Congenital heart diseases
 - 1.4.5 Schistosomiasis
- 1.5 PAH long-term responders to calcium channel blockers
- 1.6 PAH with overt features of venous/capillaries (PVOD/PCH) involvement
- 1.7 Persistent PH of the newborn syndrome

2 PH due to left heart disease

- 2.1 PH due to heart failure with preserved left ventricular ejection fraction (LVEF)
- 2.2 PH due to heart failure with reduced LVEF
- 2.3 Valvular heart disease
- 2.4 Congenital/acquired cardiovascular conditions leading to post-capillary PH

3 PH due to lung diseases and/or hypoxia

- 3.1 Obstructive lung disease
- 3.2 Restrictive lung disease
- 3.3 Other lung disease with mixed restrictive/obstructive pattern
- 3.4 Hypoxia without lung disease
- 3.5 Developmental lung disorders

4 PH due to pulmonary artery obstruction

- 4.1 Chronic thromboembolic PH

Reference number(s)
1651-A

- 4.2 Other pulmonary artery obstructions
 - 4.2.1 Sarcoma (high or intermediate grade) or angiosarcoma
 - 4.2.2 Other malignant tumors
 - Renal carcinoma
 - Uterine carcinoma
 - Germ cell tumours of the testis
 - Other tumours
 - 4.2.3 Non-malignant tumours
 - Uterine leiomyoma
 - 4.2.4 Arteritis without connective tissue disease
 - 4.2.5 Congenital pulmonary artery stenosis
 - 4.2.6 Parasites
 - Hydatidosis

5 PH with unclear and/or multifactorial mechanisms

- 5.1 Hematologic disorders: Chronic hemolytic anemia, myeloproliferative disorders
- 5.2 Systemic and metabolic disorders: Pulmonary Langerhans cell histiocytosis, Gaucher disease, glycogen storage disease, neurofibromatosis, sarcoidosis
- 5.3 Others: chronic renal failure with or without hemodialysis, fibrosing mediastinitis
- 5.4 Complex congenital heart disease

VI. REFERENCES

1. Revatio [package insert]. Morgantown, WV; Viatrix Specialty LLC; January 2023.
2. Sildenafil [package insert]. Morgantown, WV: Mylan Pharmaceuticals Inc.; February 2018.
3. Sildenafil for oral suspension [package insert]. Piscataway, NJ: Novadoz Pharmaceuticals LLC; June 2020.
4. Liqrev [package insert]. Farmville, NC: CMP Pharma, Inc.; April 2023.
5. Chin KM, Rubin LJ. Pulmonary arterial hypertension. *J Am Coll Cardiol.* 2008;51(16):1527-1538.
6. McLaughlin VV, Archer SL, Badesch DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association developed in collaboration with the American College of Chest Physicians; American Thoracic Society, Inc.; and the Pulmonary Hypertension Association. *J Am Coll Cardiol.* 2009;53(17):1573-1619.
7. Badesch DB, Champion HC, Gomez-Sanchez MA, et al. Diagnosis and assessment of pulmonary arterial hypertension. *J Am Coll Cardiol.* 2009;54:S55-S66.
8. Simonneau G, Robbins IM, Beghetti M, et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol.* 2013;62:D34-S41.
9. Rubin LJ; American College of Chest Physicians. Diagnosis and management of pulmonary arterial hypertension: ACCP evidence-based clinical practice guidelines. *Chest.* 2004;126(1 Suppl):7S-10S.
10. Barst RJ, Gibbs SR, Ghofrani HA, et al. Updated evidence-based treatment algorithm in pulmonary arterial hypertension. *J Am Coll Cardiol.* 2009;54:S78-S84.
11. Taichman DB, Ornelas J, Chung L, et al. Pharmacologic therapy for pulmonary arterial hypertension in adults. CHEST guideline and expert panel report. *Chest.* 2014;46(2):449-475.
12. Barst RJ, Ivy DD, Gaitan G, et al. A randomized, double-blind, placebo-controlled, dose-ranging study of oral sildenafil citrate in treatment-naïve children with pulmonary arterial hypertension. *Circulation.* 2012;125:324-334.
13. FDA Drug Safety Communication: FDA recommends against use of Revatio in children with pulmonary hypertension. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm317123.htm>. Accessed April 28, 2015.

Reference number(s)
1651-A

14. Abman SH, Kinsella JP, Rosenzweig EB, et al. Implications of the U.S. Food and Drug Administration warning against the use of sildenafil for the treatment of pediatric pulmonary hypertension. *Am J Respir Crit Care Med.* 2013;187(6):572-575.
15. FDA Drug Safety Communication: FDA clarifies Warning about Pediatric Use of Revatio (sildenafil) for Pulmonary Arterial Hypertension. Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm390876.htm>. Accessed April 28, 2015.
16. Abman, SH, Hansmann G, Archer SL, et al. Pediatric pulmonary hypertension: guidelines from the American Heart Association and American Thoracic Society. *Circulation.* 2015;132(21):2037-99.
17. Galie N, Ghofrani HA, Torbicki A, et al. Sildenafil citrate therapy for pulmonary arterial hypertension. *N Engl J Med.* 2005;353:2148-2157.
18. Rubin LJ, Badesch DB, Fleming RT, et al. Long-term treatment with sildenafil citrate in pulmonary arterial hypertension. *Chest.* 2011;140(5):1274-1283.
19. IBM Micromedex® DRUGDEX® (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at: <https://www.micromedexsolutions.com>. Accessed February 8, 2023.
20. Hughes M, Ong VH, Anderson ME, et al. Consensus best practice pathway of the UK Scleroderma Study Group: digital vasculopathy in systemic sclerosis. *Rheumatology.* 2015;54:2015-2024.
21. Roustit M, Blaise S, Allanore Y, et al. Phosphodiesterase-5 inhibitors for the treatment of secondary Raynaud's phenomenon: systematic review and meta-analysis of randomized trials. *Ann Rheum Dis.* 2013;72(10):1696-1699.
22. Walker KM, Pope J, et al. Treatment of systemic sclerosis complications: what to use when first-line treatment fails – a consensus of systemic sclerosis experts. *Semin Arthritis Rheum.* 2012;42(1):42-55.
23. Klinger, JR., Elliott, CG, Levine, DJ, et al. Therapy for Pulmonary Arterial Hypertension in Adults: Update of the CHEST Guidelines and Expert Panel Report. *Chest.* 2019;155(3): 565-586.
24. Galie, N., McLaughlin, VV, Rubin, LJ, Simonneau, G. An overview of the 6th World Symposium on Pulmonary Hypertension. *Eur Respir J* 2019; 53: 1802148; DOI: 10.1183/13993003.02148-2018. Published 24 January 2019.
25. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J* 2019;53:1801913; doi:10.1183/13993003.01913-2018.
26. Barst RJ, Beghetti M, Pulido T, et al. STARTS-2: long-term survival with oral sildenafil monotherapy in treatment-naive pediatric pulmonary arterial hypertension. *Circulation.* 2014 May 13;129(19):1914-23.
27. Kwan WC, Shavelle DM, Laughrun DR. Pulmonary vascular resistance index: Getting the units right and why it matters. *Clin Cardiol.* 2019 Mar;42(3):334-338.