

## Omvo<sup>h</sup>® (mirikizumab-mrkz) (Intravenous and Subcutaneous)

Effective Date: 05/01/2024

Review Date: 02/08/2024, 07/17/2024, 1/22/2025

Pharmacy Scope (subcutaneous formulation only): Medicaid

Medical Scope (intravenous formulation only): Medicaid, Commercial, Medicare-Medicaid Plan (MMP)

### I. Length of Authorization

- **Medical Scope:**

- Coverage for intravenous (IV) Omvoh will be provided once for 12 weeks (for 3 IV doses) and may not be renewed.

\*\* For members that meet criteria, Omvoh 200 mg (subcutaneous dose) will be approved for every 4 weeks thereafter for 4 months for Medicaid and Commercial ONLY\*\*

- **Pharmacy Scope:**

- Coverage for subcutaneous (SC) Omvoh will be provided for 6 months and may be renewed for 6 months.

### II. Dosing Limits

A. **Medical Scope: Intravenous**

- **Quantity Limit (max daily dose) [NDC Unit]:**
  - Omvoh 300 mg/15 mL single-dose vial: 1 vial at Weeks 0, 4 & 8 (3 vials total)
- **Max Units (per dose and over time) [HCPS Unit]:**
  - 900 mg or 900 units per 90 days

B. **Pharmacy Scope: Subcutaneous**

- **Quantity Limit (max daily dose) [NDC Unit]:**
  - Omvoh 100 mg/mL pen injection: 2 pens per 28 days (daily dose of 0.072)

### III. Initial Approval Criteria

Coverage is provided in the following conditions:

**For all indications:**

- Submission of the member's chart or medical record is required, documenting medical necessity based on the criteria corresponding to the applicable indication; **AND**

- Member has a pretreatment tuberculosis (TB) screening with a TB skin test or an interferon gamma release assay (e.g., QFT-GIT, T-SPOT.TB). *[Note: Members who have received another biologic DMARD or targeted synthetic DMARD (e.g., Xeljanz) are exempt from requirements related to TB screening in this Policy.]; AND*
- Member is free of any clinically important active infection, including clinically important localized infections; **AND**
- Member will not receive live vaccines during therapy; **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Baseline liver enzymes and bilirubin levels have been obtained prior to initiating therapy; **AND**
- Omvoh will not be used concomitantly with an injectable biologic response modifier including TNF-inhibitors (e.g., Humira (adalimumab), Enbrel (etanercept), Remicade (infliximab), Simponi (golimumab), etc.) and IL-inhibitors (e.g., Cosentyx (secukinumab), ustekinumab (e.g., Stelara, Wezlana, etc.) ( Tremfya (guselkumab), Ilumya (tildrakizumab), Skyrizi (risankizumab), etc.) or other oral non-biologic agent (e.g., Otezla (apremilast), Xeljanz (tofacitinib), Olumiant (baricitinib), Rinvoq (upadacitinib), Velsipity (etrasimod), etc.)

#### **A. Moderately to severely active ulcerative colitis (UC)**

Authorization may be granted for treatment of moderately to severely active UC when all of the following criteria are met:

- Member is 18 years of age or older; **AND**
- This medication must be prescribed by or in consultation with a gastroenterologist; **AND**
- Documented moderate to severe UC; **AND**
- Member has had an inadequate response, intolerance, or contraindication to at least a 3-month trial of infliximab IV or adalimumab at maximum tolerated doses; **AND**
- Member has had an inadequate response, intolerance, or contraindication to at least a 6-month trial of ustekinumab biosimilar at maximum tolerated doses

## **IV. Renewal Criteria**

Authorization of 6 months may be granted for all members (including new members) when all of the following criteria are met:

- Member continues to meet all initial authorization criteria; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: anaphylaxis or other serious allergic reactions, severe infections, jaundice or other evidence of significant liver injury, etc.; **AND**
- Member has annual eye exams to monitor for macular edema; **AND**
- Member is using the requested medication for moderate to severe active ulcerative colitis and one of the following must apply:
  - Member has achieved or maintained remission; **OR**
  - Member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
    - a. Stool frequency

- b. Rectal bleeding
- c. Urgency of defecation
- d. C-reactive protein (CRP)
- e. Fecal calprotectin (FC)
- f. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
- g. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

## V. Dosage/Administration

Indication	Dose
<b>Moderately to severely active ulcerative colitis (UC)</b>	<ul style="list-style-type: none"> <li>IV- Induction dosage is 300 mg administered by intravenous infusion over at least 30 minutes at Weeks 0, 4, and 8.</li> <li>SC- Maintenance dosage is 200 mg administered by subcutaneous injection (given as two consecutive injections of 100 mg each) at Week 12, and every 4 weeks thereafter.</li> </ul>
<ul style="list-style-type: none"> <li>The vial and prefilled pen are not made with dry natural rubber latex.</li> <li>OMVOH (mirikizumab-mrkz) injection is a sterile, preservative-free, clear to opalescent, colorless to slightly yellow to slightly brown solution for intravenous infusion or subcutaneous injection.</li> <li>Each single-dose prefilled pen consists of a 1 mL glass syringe with a fixed 27-gauge ½ inch needle.</li> </ul>	

## VI. Billing Code/Availability Information

### HCPCS:

- J2267 – Injection, mirikizumab-mrkz, 1 mg

### NDC:

- OMVOH (mirikizumab-mrkz) injection is a sterile, preservative-free, clear to opalescent, colorless to slightly yellow to slightly brown solution for intravenous infusion or subcutaneous injection.: 0002-7575-01

## VII. References

1. Omvoh [package insert]. Indianapolis, IN: Eli Lilly and Company; October 2023.