

Evolent Clinical Guideline 3027 for Zaltrap[™] (zivaflibercept)

Guideline Number: Evolent_CG_3027	Applicable Codes		
"Evolent" refers to Evolent Health LLC and Evolent Specialty Services, Inc. © 2012 - 2025 Evolent. All rights Reserved.			
Original Date: October 2012	Last Revised Date: February 2025	Implementation Date: February 2025	

TABLE OF CONTENTS

STATEMENT	2
Purpose	2
INDICATIONS	2
COLORECTAL CANCER	2
CONTRAINDICATIONS/WARNINGS	2
EXCLUSION CRITERIA	2
CODING AND STANDARDS	3
CODES	3
APPLICABLE LINES OF BUSINESS	3
POLICY HISTORY	4
LEGAL AND COMPLIANCE	Δ
GUIDELINE APPROVAL	
Committee	4
DISCLAIMER	4
REFERENCES	5



STATEMENT

Purpose

To define and describe the accepted indications for Zaltrap (ziv-aflibercept) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

INDICATIONS

Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided

- The member has not experienced disease progression on the requested medication AND
- The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization AND
- Additional medication(s) are not being added to the continuation request.

Colorectal Cancer

NOTE: Zaltrap (ziv-aflibercept) is not supported by Evolent Policy for metastatic colorectal cancer. This policy position is based on the lack of Level 1 Evidence (randomized clinical trials and/or meta-analyses) to show superior outcomes with Zaltrap (ziv-aflibercept) based regimens compared to Avastin (bevacizumab)/bevacizumab biosimilars based regimens. Furthermore, meta-analyses have shown an increased risk of infections and proteinuria with this agent, please see reference below. Please refer to alternative agents/regimens recommended by Evolent, including but not limited to regimens available at <u>Evolent Pathways</u>

CONTRAINDICATIONS/WARNINGS

None

EXCLUSION CRITERIA

- Disease progression while taking Zaltrap (ziv-aflibercept).
- Dosing exceeds single dose limit of Zaltrap (ziv-aflibercept) 4 mg/kg.

Page 2 of 5



- Investigational use of Zaltrap (ziv-aflibercept) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community.
 Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 - Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 - Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 - o Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 - Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 - That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
 - That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
 - That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

CODING AND STANDARDS

Codes

• J9400 - Injection, ziv-aflibercept, 1 mg

Applicable Lines of Business

	CHIP (Children's Health Insurance Program)
\boxtimes	Commercial
\boxtimes	Exchange/Marketplace
\boxtimes	Medicaid
	Medicare Advantage



POLICY HISTORY

Date	Summary
February 2025	 Converted to new Evolent guideline template This guideline replaces UM ONC_1226 Zaltrap (zivaflibercept)
February 2024	Updated NCH verbiage to EvolentUpdated "continuation request" verbiage

LEGAL AND COMPLIANCE

Guideline Approval

Committee

Reviewed / Approved by Evolent Specialty Clinical Guideline Review Committee

Disclaimer

Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment, and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. Members should contact their Plan customer service representative for specific coverage information.



REFERENCES

- Van Cutsem E, et al. Addition of aflibercept to fluorouracil, leucovorin, and irinotecan improves survival in a phase III randomized trial in patients with metastatic colorectal cancer previously treated with an oxaliplatin-based regimen. *J Clin Oncol*. 2012 Oct 1;30(28):3499-506. doi: 10.1200/JCO.2012.42.8201.
- 2. Zhang X, et al. Incidence and risk of severe infections associated with aflibercept in cancer patients: a systematic review and meta-analysis. *Br J Clin Pharmacol*. 2016 Jan;81(1):33-40. doi: 10.1111/bcp.12758.
- 3. Zaltrap prescribing information. Sanofi-Aventis U.S. LLC, Bridgewater, NJ 2023.
- AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs Bethesda, MD 2025.
- 5. Clinical Pharmacology Elsevier Gold Standard 2025.
- 6. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2025.
- 7. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2025.
- 8. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol*. 2014 Apr 20;32(12):1277-80.
- Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf.