

Evolut Clinical Guideline 3122 for Monjuvi™ (tafasitamab-cxix)

Guideline Number: Evolut_CG_3122	<u>Applicable Codes</u>	
<i>"Evolut" refers to Evolut Health LLC and Evolut Specialty Services, Inc.</i> <i>© 2020 - 2025 Evolut. All rights Reserved.</i>		
Original Date: September 2020	Last Revised Date: August 2025	Implementation Date: August 2025

TABLE OF CONTENTS

STATEMENT	2
PURPOSE	2
INDICATIONS	2
DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL)	2
FOLLICULAR LYMPHOMA (FL)	2
CONTRAINDICATIONS/WARNINGS	3
EXCLUSION CRITERIA	3
CODING AND STANDARDS	4
CODES	4
APPLICABLE LINES OF BUSINESS	4
POLICY HISTORY	4
LEGAL AND COMPLIANCE	5
GUIDELINE APPROVAL	5
Committee	5
DISCLAIMER	5
REFERENCES	5

STATEMENT

Purpose

To define and describe the accepted indications for Monjuvi (tafasitamab-cxix) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

INDICATIONS

Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided

- The member has not experienced disease progression on the requested medication AND
- The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization AND
- Additional medication(s) are not being added to the continuation request.

Diffuse Large B-Cell Lymphoma (DLBCL)

- Monjuvi (tafasitamab-cxix) may be used in combination with Revlimid (lenalidomide) for up to 12 cycles with further continuation as monotherapy for members with all of the following:
 - CD-19 expression on lymphoma cells confirmed by any standard test AND
 - Relapsed/refractory DLBCL (including transformation from an indolent lymphoma) who are not candidates for a stem cell transplant or have failed a prior stem cell transplant AND
 - Receipt of 1-3 prior regimens including a chemoimmunotherapy regimen containing an anti-CD20-targeted therapy (e.g., rituximab, obinutuzumab).

Follicular Lymphoma (FL)

- Monjuvi (tafasitamab-cxix) may be used in combination with Revlimid (lenalidomide) and rituximab/rituximab biosimilar in adult members with relapsed or refractory follicular lymphoma (FL) who have failed at least one systemic therapy, including an anti-CD20 monoclonal antibody (e.g., rituximab/rituximab biosimilar, obinutuzumab).

CONTRAINDICATIONS/WARNINGS

- None

EXCLUSION CRITERIA

- Monjuvi (tafasitamab-cxix) is being used on or after disease progression with the same regimen
- Lack of confirmed documentation of CD19 expression on lymphoma cells confirmed by any standard test.
- Dosing exceeds single dose limit of Monjuvi (tafasitamab-cxix) 12 mg/kg.
- Treatment exceeds the maximum duration limit of 12 cycles (when used in combination with lenalidomide).
- Investigational use of Monjuvi (tafasitamab-cxix) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 - Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 - Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 - Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 - Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 - That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
 - That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
 - That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

CODING AND STANDARDS

Codes

- J9349 - Injection, tafasitamab-cxix, 2 mg

Applicable Lines of Business

☐	CHIP (Children's Health Insurance Program)
☒	Commercial
☒	Exchange/Marketplace
☒	Medicaid
☐	Medicare Advantage

POLICY HISTORY

Date	Summary
August 2025	• Updated indication section with Follicular Lymphoma indication • Updated references
June 2025	• Converted to new Evolent guideline template • This guideline replaces UM ONC_1412 Monjuvi (tafasitamab-cxix) • Updated indication section • Updated exclusion criteria • Updated references
June 2024	• Updated NCH verbiage to Evolent

LEGAL AND COMPLIANCE

Guideline Approval

Committee

Reviewed / Approved by Evolent Specialty Clinical Guideline Review Committee

Disclaimer

Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment, and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. Evolent clinical guidelines contain guidance that requires prior authorization and service limitations. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. Members should contact their Plan customer service representative for specific coverage information.

REFERENCES

1. Sehn LH, et al. Tafasitamab Plus Lenalidomide and Rituximab for Relapsed or Refractory Follicular Lymphoma: Results from a Phase 3 Study (inMIND). *Blood*. 2024 November 25;144(2): LBA-1-LBA-4. <https://doi.org/10.1182/blood-2024-212970>.
2. Salles G, et al. Tafasitamab plus lenalidomide in relapsed or refractory diffuse large B-cell lymphoma (L-MIND): a multicentre, prospective, single-arm, phase 2 study. *Lancet Oncol*. 2020 Jul;21(7):978-988. doi: 10.1016/S1470-2045(20)30225-4.
3. Duell J, et al. Long-term outcomes from the Phase II L-MIND study of tafasitamab (MOR208) plus lenalidomide in patients with relapsed or refractory diffuse large B-cell lymphoma. *Haematologica*. 2021 Sep 1;106(9):2417-2426. doi: 10.3324/haematol.2020.275958.
4. Nowakowski GS, et al. Improved Efficacy of Tafasitamab plus Lenalidomide versus Systemic Therapies for Relapsed/Refractory DLBCL: RE-MIND2, an Observational Retrospective Matched Cohort Study. *Clin Cancer Res*. 2022 Sep 15;28(18):4003-4017. doi: 10.1158/1078-0432.CCR-21-3648.
5. Monjuvi prescribing information. Incyte Corporation. Wilmington, DE 2025.
6. Clinical Pharmacology Elsevier Gold Standard 2025.
7. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2025.
8. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2025.
9. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs. Bethesda, MD 2025.
10. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol*. 2014 Apr 20;32(12):1277-80.

11. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services:
<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
12. Current and Resolved Drug Shortages and Discontinuations Reported to the FDA:
<http://www.accessdata.fda.gov/scripts/drugshortages/default.cfm>.