

Evolut Clinical Guideline 3165 for Treanda/Bendeka/Belrapzo/Vivimusta™ (bendamustine)

Guideline Number: Evolut_CG_3165	<u>Applicable Codes</u>	
<i>"Evolut" refers to Evolut Health LLC and Evolut Specialty Services, Inc.</i> © 2012 - 2025 Evolut. All rights Reserved.		
Original Date: September 2012	Last Revised Date: August 2025	Implementation Date: August 2025

TABLE OF CONTENTS

STATEMENT	2
PURPOSE	2
INDICATIONS	2
CHRONIC LYMPHOCYTIC LEUKEMIA/SMALL LYMPHOCYTIC LYMPHOMA (CLL/SLL)	2
NON-HODGKIN'S LYMPHOMA (NHL)	2
SMALL CELL LUNG CANCER (SCLC)	3
CONTRAINDICATIONS/WARNINGS	3
EXCLUSION CRITERIA	3
CODING AND STANDARDS	4
CODES	4
APPLICABLE LINES OF BUSINESS	4
POLICY HISTORY	5
LEGAL AND COMPLIANCE	5
GUIDELINE APPROVAL	5
Committee	5
DISCLAIMER	5
REFERENCES	5

STATEMENT

Purpose

To define and describe the accepted indications for Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

Evolent is responsible for processing all medication requests from network ordering providers. Medications not authorized by Evolent may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

INDICATIONS

Continuation requests for a not-approvable medication shall be exempt from this Evolent policy provided

- The member has not experienced disease progression on the requested medication AND
- The requested medication was used within the last year without a lapse of more than 30 days of having an active authorization AND
- Additional medication(s) are not being added to the continuation request.

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL)

- Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used in combination with rituximab/rituximab biosimilar as initial or subsequent therapy for members with CLL.

Non-Hodgkin's Lymphoma (NHL)

- Indolent B-Cell Lymphomas: Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used in the following clinical settings:
 - In combination with rituximab/rituximab biosimilar for primary or subsequent therapy of any of the following:
 - Follicular B-Cell Lymphoma
 - Nodal Marginal Zone/Extra-Nodal Marginal Zone/Gastric MALT Lymphoma/Non-Gastric MALT Lymphoma/Splenic Marginal Zone Lymphoma.
- Diffuse Large B-Cell Lymphoma
 - Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used as subsequent therapy for relapsed or refractory disease in combination with Polivy (polatuzumab) with or without rituximab/rituximab biosimilar.

- Mantle Cell Lymphoma
 - Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) may be used as initial or subsequent therapy in combination with rituximab/rituximab biosimilar.

Small Cell Lung Cancer (SCLC)

- NOTE: Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) products are not supported by Evolent Policy for the treatment of relapsed/refractory SCLC. This policy position is based on the lack of Level 1 Evidence (randomized clinical trials and/or meta-analyses) to show superior outcomes compared to Evolent recommended regimens, available at [**Evolent Pathways**](#).

CONTRAINDICATIONS/WARNINGS

- Contraindications
 - Treanda/Bendeka/Belrapzo/Vivimusta
 - Hypersensitivity (e.g., anaphylactic or anaphylactoid reactions) to bendamustine or any component of the formulation
 - Bendeka/Belrapzo
 - Hypersensitivity to polyethylene glycol 400, propylene glycol, or monothioglycerol
 - Vivimusta
 - Hypersensitivity to polyethylene glycol 400, dehydrated alcohol, or monothioglycerol

EXCLUSION CRITERIA

- Member has disease progression while on Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine).
- Dosing exceeds single dose limit of Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) 100 mg/m² for CLL and 120 mg/m² for NHL.
- Treatment with Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) exceeds the maximum duration limit of 8 cycles for NHL and 6 cycles for CLL.
- Investigational use of Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 - Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 - Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 - Whether the reported study outcomes represent clinically meaningful outcomes

experienced by patients. Generally, the definitions of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.

- Whether the experimental design, considering the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
- That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
- That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
- That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

CODING AND STANDARDS

Codes

- J9033 - Injection, bendamustine hydrochloride, 1 mg
- J9034 - Injection, bendamustine hcl (bendeka), 1 mg
- J9036 - Injection, bendamustine hydrochloride, (belrapzo/bendamustine), 1 mg
- J9056 - Injection, bendamustine hydrochloride (vivimusta), 1 mg

Applicable Lines of Business

☐	CHIP (Children's Health Insurance Program)
☒	Commercial
☒	Exchange/Marketplace
☒	Medicaid
☐	Medicare Advantage

POLICY HISTORY

Date	Summary
August 2025	- Converted to new Evolent guideline template - This guideline replaces UM ONC_1215 Treanda/Bendeka/Belrapzo/Vivimusta (bendamustine) - Updated indication section - Updated references
August 2024	- Added new product “Vivimusta” to policy - Added new reference - Updated NCH verbiage to Evolent

LEGAL AND COMPLIANCE

Guideline Approval

Committee

Reviewed / Approved by Evolent Specialty Clinical Guideline Review Committee

Disclaimer

Evolent Clinical Guidelines do not constitute medical advice. Treating health care professionals are solely responsible for diagnosis, treatment, and medical advice. Evolent uses Clinical Guidelines in accordance with its contractual obligations to provide utilization management. Coverage for services varies for individual members according to the terms of their health care coverage or government program. Individual members' health care coverage may not utilize some Evolent Clinical Guidelines. Evolent clinical guidelines contain guidance that requires prior authorization and service limitations. A list of procedure codes, services or drugs may not be all inclusive and does not imply that a service or drug is a covered or non-covered service or drug. Evolent reserves the right to review and update this Clinical Guideline in its sole discretion. Notice of any changes shall be provided as required by applicable provider agreements and laws or regulations. Members should contact their Plan customer service representative for specific coverage information.

REFERENCES

1. Treanda prescribing information. Teva Pharmaceuticals. Parsippany, NJ 2022.
2. Vivimusta prescribing information. Slayback Pharma LLC. Princeton, NJ 2024.
3. Bendeka prescribing information. Teva Pharmaceuticals. Parsippany, NJ 2024.
4. Belrapzo prescribing information. Eagle Pharmaceuticals, Inc. Woodcliff Lake, NJ. 2024.
5. Clinical Pharmacology Elsevier Gold Standard 2025.

6. Micromedex® Healthcare Series: Micromedex Drugdex Ann Arbor, Michigan 2025.
7. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2025.
8. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs. Bethesda, MD 2025.
9. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol*. 2014 Apr 20;32(12):1277-80.
10. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services:
<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
11. Current and Resolved Drug Shortages and Discontinuations Reported to the FDA:
<http://www.accessdata.fda.gov/scripts/drugshortages/default.cfm>.