

Evenity™ (romosozumab-aqqg) (Subcutaneous)

Effective date:10/01/2019

Review Date: 6/1/2020, 5/20/2021, 3/3/2022, 3/23/2023, 12/14/2023, 01/04/2024, 04/30/2025, 02/03/2026

Scope: Medicaid, Commercial, Medicare

I. Length of Authorization

Coverage will be provided for 12 months and may NOT be renewed.

II. Dosing Limits

A. Max Units (per dose and over time) [HCPCS Unit]:

- 210 billable units every month

III. Summary of Evidence

Evenity is a sclerostin inhibitor indicated for the treatment of osteoporosis in postmenopausal women at high risk for fracture, defined as a history of osteoporotic fracture, or multiple risk factors for fracture; or patients who have failed or are intolerant to other available osteoporosis therapy. Clinical studies show that Evenity produces rapid increases in bone formation within two weeks of injection and maintains therapeutic activity for several months, supporting significant improvements in bone mineral density and fracture risk reduction. The most common adverse effects include joint pain (arthralgia), headache, and injection-site irritation.

IV. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

Medicare members who have previously received this medication within the past 365 days are not subject to Step Therapy Requirements

- Member is at least 18 years of age; **AND**
- Confirmation member is receiving calcium and Vitamin D supplementation if dietary intake is inadequate; **AND**
- Member must not have hypocalcemia; **AND**
- Member has not had a myocardial infarction or stroke within the preceding year (*Note: in members with other cardiovascular disease and/ or risk factors, consider whether benefits of therapy outweigh the risks.*); **AND**

Osteoporosis in Women † ¹⁻¹²

- Member must be at a high risk for fracture**; **AND**
- Member must be post-menopausal; **AND**
- Member has a documented diagnosis of osteoporosis indicated by one or more of the following:

- Hip/femur DXA (femoral neck or total hip) or lumbar spine T-score ≤ -2.5 and/or forearm DXA at the 33% (one-third) radius site; **OR**
- T-score ≤ -1 or low bone mass and a history of fragility fracture to the hip or spine; **OR**
- T-score between -1 and -2.5 with a FRAX 10-year probability for major fracture $\geq 20\%$ or hip fracture $\geq 3\%$; **AND**
- § Documented treatment failure or ineffective response[±] to a minimum (12) month trial on previous therapy with bisphosphonates (oral or IV) such as alendronate, risedronate, ibandronate, or zoledronic acid; **OR**
 - Member has a documented contraindication* or intolerance to BOTH oral bisphosphonates AND intravenous (IV) bisphosphonates such as alendronate, risedronate, ibandronate, or zoledronic acid; **AND**
- § Documented treatment failure or ineffective response[±] to a minimum (12) month trial on previous therapy with RANKL-blocking agents such as denosumab, etc.; **OR**
 - Member has a documented contraindication* or intolerance to RANKL-blocking agents such as denosumab, etc.

§ Members with very high risk for fracture defined as a T-score ≤ -3.0 , or a T-score ≤ -2.5 with a history of fragility fractures or severe or multiple vertebral fractures are not subject to prior trial and failure requirements with bisphosphonates and/or denosumab^{9,10,11}

± Ineffective response is defined as one or more of the following:⁵

- Decrease in T-score in comparison with baseline T-score from DXA scan
- Patient has a new fracture while on bisphosphonate therapy

**** High risk for fractures include, but are not limited to, one or more of the following:⁵**

- History of an osteoporotic fracture as an adult
- Parental history of hip fracture
- Low BMI
- Rheumatoid arthritis
- Alcohol intake (3 or more drinks per day)
- Current smoking
- History of oral glucocorticoids ≥ 5 mg/d of prednisone (or equivalent) for >3 months (ever)

*** Examples of contraindications to oral bisphosphonate therapy include the following:**

- Documented inability to sit or stand upright for at least 30 minutes
- Documented pre-existing gastrointestinal disorder such as inability to swallow, Barrett's esophagus, esophageal stricture, dysmotility, or achalasia
- Surgical anastomoses are present in the GI tract after certain types of bariatric surgery (e.g., Roux-en-Y gastric bypass).

*** Examples of contraindications to injectable bisphosphonate therapy include the following:**

- Documented pre-existing hypocalcemia and disturbances of mineral metabolism
- Documented pre-existing renal insufficiency defined as creatinine clearance < 35 mL/min

*** Examples of contraindications to RANKL-blocking therapy include the following:**

- Documented pre-existing hypocalcemia and disturbances of mineral metabolism
- Documented hypersensitivity to the active ingredient or its excipients

† FDA Approved Indication(s); ‡ Compendia recommended indication(s)

V. Renewal Criteria

Coverage may NOT be renewed.

VI. Dosage/Administration

Indication	Dose
Osteoporosis	Administer 210 mg subcutaneously (as two separate subcutaneous injections of 105 mg each) by a health care provider every month for a total of 12* monthly doses.

**Note: The anabolic effect of Evenity wanes after 12 monthly doses of therapy. Therefore, the duration of Evenity use should be limited to 12 monthly doses. If osteoporosis therapy remains warranted, continued therapy with an anti-resorptive agent should be considered.*

VII. Billing Code/Availability Information

HCPCS Code:

- J3111 – Injection, romosozumab-aqqg, 1 mg; 1 billable unit = 1 mg

NDC:

- Evenity 105 mg/1.17 mL single-use prefilled syringe: 55513-0880-xx

VIII. References

1. Evenity [package insert]. Thousand Oaks, CA; Amgen, Inc.; January 2026. Accessed January 2026.
2. Cosman F, Crittenden DB, Ferrari S, Khan A, Lane NE, Lippuner K, Matsumoto T, Milmont CE, Libanati C, Grauer A. FRAME Study: The Foundation Effect of Building Bone With 1 Year of Romosozumab Leads to Continued Lower Fracture Risk After Transition to Denosumab. *J Bone Miner Res.* 2018 Jul;33(7):1219-1226. doi: 10.1002/jbmr.3427. Epub 2018 May 17.
3. WHO Scientific Group on the Prevention and Management of Osteoporosis. Prevention and management of osteoporosis: report of a WHO scientific group. (WHO technical report series; 921). Geneva, Switzerland: WHO; 2000.
4. Kanis JA on behalf of the World Health Organization Scientific Group (2007). Assessment of osteoporosis at the primary health care level. Technical Report. World Health Organization Collaborating Center for Metabolic Bone Diseases. University of Sheffield, UK; 2007.
5. National Osteoporosis Foundation. Clinician's Guide to Prevention and Treatment of Osteoporosis. Washington, DC: National Osteoporosis Foundation; 2014.

6. Camacho PM, Petak SM, Binkley N, et al. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS AND AMERICAN COLLEGE OF ENDOCRINOLOGY CLINICAL PRACTICE GUIDELINES FOR THE DIAGNOSIS AND TREATMENT OF POSTMENOPAUSAL OSTEOPOROSIS - 2016. *Endocr Pract.* 2016 Sep 2; 22(Suppl 4):1-42.
7. Qaseem A, Forcica MA, McLean RM, Denberg TD; Clinical Guidelines Committee of the American College of Physicians. Treatment of Low Bone Density or Osteoporosis to Prevent Fractures in Men and Women: A Clinical Practice Guideline Update from the American College of Physicians. *Ann Intern Med.* 2017 May 9. doi: 10.7326/M15-1361.
8. Jeremiah MP, Unwin BK, Greenawald MH, et al. Diagnosis and Management of Osteoporosis. *Am Fam Physician.* 2015 Aug 15;92(4):261-8.
9. Eastell R, Rosen CJ, Black DM, et al. Pharmacological Management of Osteoporosis in Postmenopausal Women: An Endocrine Society Clinical Practice Guideline. *J Clin Endo Metab*, Vol 104, Iss 5, May 2019, pps 1595–1622, <https://doi.org/10.1210/jc.2019-00221>
10. Shoback D, Rosen CJ, Black DM, et al. Pharmacological Management of Osteoporosis in Postmenopausal Women: An Endocrine Society Guideline Update. *J Clin Endocrinol Metab*, March 2020, 105(3):1–8. <https://academic.oup.com/jcem>
11. Rosen, HN, Drezner MK. Overview of the management of osteoporosis in postmenopausal women. In: Rosen CJ, Schmader KE, ed. *UpToDate*. Waltham, MA.: UpToDate; 2019. www.uptodate.com. Accessed June 1, 2019.
12. Jackson RD, LaCroix AZ, Gass M, Women’s Health Initiative Investigators. Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med.* 2006; 354(7):669–683.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
M80.00XA- M80.08XS	Age-related osteoporosis with current pathological fracture
M81.0	Age-related osteoporosis without current pathological fracture

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs) and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

Policy Rationale:

Evenity was reviewed by the Neighborhood Health Plan of Rhode Island Pharmacy & Therapeutics (P&T) Committee. Neighborhood adopted the following clinical coverage criteria to ensure that its members use Evenity according to Food and Drug Administration (FDA) approved labeling and/or relevant clinical literature. Neighborhood worked with network prescribers and pharmacists to draft these criteria. These criteria will help ensure its members are using this drug for a medically accepted indication, while minimizing the risk for adverse effects and ensuring more cost-effective options are used first, if applicable and appropriate. For Medicare members, these coverage criteria will only apply in the absence of National Coverage Determination (NCD) or Local Coverage Determination (LCD) criteria. Neighborhood will give individual consideration to each request it reviews based on the information submitted by the prescriber and other information available to the plan.